

# PHY411/AST233 Lecture notes -Introduction to Hamiltonian and Lagrangian mechanics

Alice Quillen

November 4, 2024

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
1.1	Hamiltonian Mechanics . . . . .	2
1.2	Lagrangian mechanics . . . . .	6
1.3	Dynamical systems . . . . .	6
1.4	Go down a dimension . . . . .	6
1.5	Even more abstractly - Symbolic dynamics . . . . .	7
1.6	Infinite dimensional systems . . . . .	7
<b>2</b>	<b>Equations of motion with one degree of freedom</b>	<b>8</b>
2.1	Energy is conserved . . . . .	8
2.2	Phase flow . . . . .	9
2.2.1	Example: The Harmonic oscillator . . . . .	9
2.2.2	Trajectories can go in either direction . . . . .	10
2.2.3	What is a fixed point? . . . . .	10
2.2.4	Equivalent Hamiltonian model . . . . .	10
2.2.5	Example: Pendulum . . . . .	13
2.2.6	Example: Cubic Potential . . . . .	14
2.2.7	Example: The radial degree of freedom in Kepler's problem . . . . .	16
2.3	Period as a function of area in phase space . . . . .	18
2.3.1	Harmonic and an-harmonic oscillator examples . . . . .	18
2.3.2	Hamiltonians that give the same orbital trajectories . . . . .	19
2.4	The separatrix of a pendulum . . . . .	20
2.5	On Hamiltonian systems with one degree of freedom . . . . .	20

<b>3</b>	<b>Relation between Lagrangian and Hamiltonian Descriptions</b>	<b>21</b>
3.1	Extremum of a path . . . . .	21
3.2	Lagrangian's equations for a Newtonian system . . . . .	22
3.3	Hamilton's equations for a Newtonian system . . . . .	22
3.4	Legendre Transforms . . . . .	24
3.4.1	Example: Legendre transform of sqrt(x) . . . . .	25
3.5	Constructing a Hamiltonian from a Lagrangian . . . . .	26
3.5.1	Example: Hamiltonian for a Newtonian system in one dimension . .	27
3.6	Geometrical view of a Lagrangian path integral . . . . .	27
3.7	Pathological Legendre Transformations . . . . .	28
3.8	Phase Space Action . . . . .	31
<b>4</b>	<b>Stability of fixed points</b>	<b>31</b>
4.1	Lyapunov and asymptotic stability . . . . .	34
<b>5</b>	<b>Dimensional rescaling for Hamiltonians</b>	<b>34</b>
<b>6</b>	<b>Liouville's Theorem and Phase wrapping</b>	<b>37</b>
6.1	The meaning of incompressible flow . . . . .	38
6.2	Phase wrapping . . . . .	39
<b>7</b>	<b>Connections between classical and quantum systems</b>	<b>42</b>
7.1	Path integrals . . . . .	42
7.2	Commutation relations . . . . .	43
7.3	Quantum chaos . . . . .	44
7.4	Decoherence . . . . .	44
7.5	Other phenomena . . . . .	44
<b>8</b>	<b>Some non-conservative two-dimensional dynamical systems</b>	<b>45</b>
8.1	Example: Damped pendulum . . . . .	45
8.1.1	Fixed Points . . . . .	46
8.1.2	Appearance of a limit cycle . . . . .	47

# 1 Introduction

## 1.1 Hamiltonian Mechanics

In this formalism the dynamics is described with a physically motivated function, known as a Hamiltonian, that depends on two types of variables, coordinates  $\mathbf{q}$ , and momenta  $\mathbf{p}$ , that are vectors of the same dimension;

$$H(\mathbf{p}, \mathbf{q}).$$

The Hamiltonian could also depend on time  $t$ , as in  $H(\mathbf{q}, \mathbf{p}, t)$ . A Hamiltonian obeys *Hamilton's equations*

$$\dot{\mathbf{q}} = \frac{\partial H}{\partial \mathbf{p}} \quad \dot{\mathbf{p}} = -\frac{\partial H}{\partial \mathbf{q}}. \quad (1)$$

Here  $\dot{\mathbf{q}} = \frac{d\mathbf{q}}{dt}$ . The difference between momenta  $\mathbf{p}$  and coordinates  $\mathbf{q}$  is that the time derivative for momenta contains a minus sign. Hamilton's equations treat coordinates and momenta differently. Let's be specific by what we mean by equation 1. Suppose  $H(x, y, p_x, p_y)$ , then Hamilton's equations are

$$\begin{aligned} \frac{dx}{dt} &= \frac{\partial H}{\partial p_x} & \frac{dp_x}{dt} &= -\frac{\partial H}{\partial x} \\ \frac{dy}{dt} &= \frac{\partial H}{\partial p_y} & \frac{dp_y}{dt} &= -\frac{\partial H}{\partial y}. \end{aligned}$$

Let's look for the moment at the lowest dimensional Hamiltonian system We can consider the dynamics of the system with a 2dimensional vector  $\mathbf{x} = (q, p)$ . This 2-dimensional space is called *phase space*. Our Hamiltonian is  $H(\mathbf{x})$ . Hamilton's equations involve partial derivatives of  $H$  or the gradient operator

$$\nabla H = \left( \frac{\partial H}{\partial q}, \frac{\partial H}{\partial p} \right).$$

Suppose we think of the Hamiltonian as an energy and insist that energy is conserved as a trajectory moves in *phase space*. For energy to be conserved, we require that the direction of motion  $\dot{\mathbf{x}}$  lies along level energy curves, as shown in Figure 1. In other words, the direction of motion (like a velocity) should be parallel to a curve of constant  $H$ . This implies that  $\dot{\mathbf{x}}$  is **perpendicular** to the gradient  $\nabla H$ ;

$$\dot{\mathbf{x}} \perp \nabla H. \quad (2)$$

This is equivalent to

$$(\mathbf{q} \ \mathbf{p}) \cdot \begin{pmatrix} \frac{\partial H}{\partial \mathbf{q}} \\ \frac{\partial H}{\partial \mathbf{p}} \end{pmatrix} = 0. \quad (3)$$

Hamilton's equations (equation 1) automatically satisfy this condition. The constant of proportionality, determining  $\dot{\mathbf{x}}$ , sets the units for time.

Hamilton's equations (equations 1 in vector form

$$\begin{pmatrix} \dot{\mathbf{q}} \\ \dot{\mathbf{p}} \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} \frac{\partial H}{\partial \mathbf{q}} \\ \frac{\partial H}{\partial \mathbf{p}} \end{pmatrix}$$




Figure 1: For energy to be conserved along a trajectory in phase space, the direction of motion should be perpendicular to the gradient of the energy function which is the Hamiltonian  $H(q, p)$ . Trajectories are level curves of the Hamiltonian and the direction of motion  $(\dot{p}, \dot{q}) \perp \nabla H$ .

can also be written as

$$\dot{\mathbf{x}} = \boldsymbol{\omega} \nabla H, \quad (4)$$

where the matrix

$$\boldsymbol{\omega} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

A symplectic  $2 \times 2$  matrix,  $g$ , is one such that

$$g^t \boldsymbol{\omega} g = \boldsymbol{\omega} \quad (5)$$

where  $g^t$  is the transpose of  $g$ . This can be generalized to higher dimensions with

$$\boldsymbol{\omega} = \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix}$$

where  $I_n$  is an  $n \times n$  identity matrix.

Hamiltonian systems are special dynamical systems in that the equations of motion generate symplectic maps of coordinates and momenta and as a consequence preserve volume in phase space. We will show what we mean by that below.

One can attempt to describe the dynamics in a coordinate independent or geometrical manner. Or using a ‘canonical’ formalism, search for nice coordinate systems that make the dynamics appear simpler.

If the Hamiltonian is only dependent on momenta

$$H(\mathbf{p})$$

then Hamilton's equations gives  $\dot{\mathbf{p}} = \frac{\partial H}{\partial \mathbf{q}} = 0$ . This implies that  $\mathbf{p} = \text{constant}$ . Hamiltonian flow preserves the momenta and we can call them **conserved quantities**. The system is said to be *integrable*. Integrable systems are interesting particularly when the dimension is high. There are examples of nonlinear system systems that are completely integrable. The dynamics of an integrable system is easy to predict. All the coordinates advance at velocities or frequencies

$$\dot{\mathbf{q}} = \nabla_p H = \frac{\partial H}{\partial \mathbf{p}}$$

(where the gradient involves derivatives of the different momenta). These are velocities (if each of the momentum  $q_i \in \mathbb{R}$ ) or frequencies (if each of the coordinates  $q_i$  is an angle) and they are all constant. One way to determine whether a finite dimensional system is integrable is to do Fourier analysis on its trajectories. If the Fourier transform only contains a few frequencies and their harmonics then the system is behaving as if it were integrable.

Suppose coordinates are real numbers and have a dimension of length  $q \sim L$ . The Hamiltonian is typically an energy, which has dimensions  $H \sim ML^2/T^2$ . If we divide the Hamiltonian by length (for computing  $\frac{\partial H}{\partial q}$ ), we have dimensions of  $ML/T^2$ . This corresponds to  $\frac{dp}{dt}$  so  $p$  must have dimension of  $ML/T$ , corresponding to momentum.

If coordinates are angles they are considered dimensionless. If the Hamiltonian is an energy per unit mass, then  $H \sim L^2/T^2$ . The momentum coordinates must have units of  $L^2/T$ . If the Hamiltonian does not depend on the angles, then the Hamiltonian is said to be in **action-angle** variables. In this case the momenta are conserved quantities called **actions**.

But there are low dimensional Hamiltonian systems that cannot be written in the form  $H(p)$ , (*chaotic ones!*). Even systems with small perturbations

$$H = H_0(\mathbf{p}) + \epsilon H_1(\mathbf{q})$$

where  $\epsilon$  is a small parameter display chaotic behavior, though if  $\epsilon$  is *small enough* then integrability is recovered. This class of systems is the focus of KAM theory.

Delineating where a complex system behaves chaotically and where it behaves like an integrable system is a challenge. There are complex looking nonlinear systems that are completely integrable (for example, the Toda lattice). And sometimes we find through numerical integration that a system behaves chaotically and would like to understand why, and predict physically relevant quantities like the rate that orbits diffuse away from an initial condition. In celestial mechanics, there can be a boundary between long lived and unstable systems that can be predicted by estimating a boundary for chaotic behavior. There can also be regions where a system only appears unstable if integrated long enough.

## 1.2 Lagrangian mechanics

In this context, the physics is dependent on a choice of Lagrangian

$$\mathcal{L}(q, \dot{q}, t)$$

A trajectory

$$\int_{t_1}^{t_2} \mathcal{L}(q, \dot{q}, t) dt.$$

is minimized in the sense that it must be on a path where the above integral is a minimum. Using calculus of variations, the system obeys Lagrange's equations

$$\frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{q}} = \frac{\partial \mathcal{L}}{\partial q}.$$

A Hamiltonian system may be transformed into a Lagrangian one and vice versa so these two approaches can be equivalent. The Lagrangian viewpoint is particularly powerful when studying constrained systems.

Exploitation of symmetries in the two formalisms is somewhat different. In Lagrangian systems we use Noether's theorem to find conserved quantities. In Hamiltonian systems we look at operators that commute with the Hamiltonian.

One can think of the Lagrangian as a way to measure *distance* between two points on a manifold or a configuration space. In this sense trajectories are *geodesics* as they minimize the distance between two points.

## 1.3 Dynamical systems

Instead of focusing on physically motivated systems, one can focus on simpler mathematical descriptions that are capable of exhibiting complex phenomena, for example,

$$\dot{x} = f(x)$$

A Hamiltonian system be written in the above way with vector  $\mathbf{x} = (\mathbf{q}, \mathbf{p})$ .

These systems can exhibit behavior that is exhibited by Hamiltonian systems, such as fixed points, bifurcations of fixed points, periodic orbits, ergodic behavior. While time independent Hamiltonian systems preserve energy, here we can also study dissipative systems. While Hamiltonian systems are area preserving, dissipative ones are not. *Strange attractors* are fractal-like features that arise in dissipative chaotic maps.

## 1.4 Go down a dimension

Starting with a Hamiltonian system we can look at the dynamics on a smaller dimensional surface, for example orbits in 4D phase space as they cross a 2D surface. This makes

*surfaces of section*. Or we can study the dynamics every orbital period or a system with a periodic perturbation. In this setting *Floquet Theory* can be helpful. In both settings maps are created from a space back to itself. The dynamical system can be viewed by looking at iteratively applied maps. To understand their properties we can study *discrete dynamical systems* or iteratively applied maps from a space on to itself. Some nice examples with complex behavior include the logistic map (a map on a line segment), circle maps, the standard map (this is a 2 d map where one dimension is the circle), and Fatou and Julia sets (of rational maps on the complex plane to itself). Integrators are discrete dynamical systems that approximate continuous ones. Separatrix maps are discrete systems that are designed to illustrate some of the dynamics of higher dimensional chaotic ones.

Billiards are even simpler dynamical systems. Given a boundary shape, an initial condition is a location on the boundary  $s$  and a trajectory angle  $\theta$ . The billiard dynamics is a map

$$\mathbf{x}_{i+1} = \mathbf{f}(\mathbf{x}_i)$$

with  $\mathbf{x}_i = (s_i, \theta_i)$  the position and angle at iteration  $i$ . It's a 2 dimensional map but based on a dynamical system, just that the trajectories are all lines between bounces that occur on the boundary. The time interval between bounces is not a constant value.

## 1.5 Even more abstractly - Symbolic dynamics

We can do down a further level of abstraction and explore what is called *symbolic dynamics* where rules are made for going between different states or members of a set. We can still get things like periodic orbits and fixed points.

Our goal here is to explore the connections between these formalisms/relations for physical systems. Visualization is very helpful in this field so we will spend some time integrating trajectories and making maps on a computer. I have created some python examples using *matplotlib* and *numpy* that can be modified for use in problem sets. One approach to making progress in understanding complex phenomena is to formulate the dynamics with toy model dynamical descriptions. Checking or understanding the behavior of these models computationally can be helpful when trying to understand or predict the behavior of a complex system. Sometimes numerical studies can give insights into what type of mathematical description might be powerful.

## 1.6 Infinite dimensional systems

We find complexity even in more abstract settings. Going the other way, Hamiltonian and Lagrangian frameworks can also be used to study infinite dimensional systems. For example, the KdV equation for shallow water waves, or hydrodynamics. In these cases, fields are a function of space and time.

## 2 Equations of motion with one degree of freedom

We consider Newton's equations

$$\ddot{x} = f(x)$$

The acceleration depends on the force  $f(x)$  that is a function of position  $x$ . The kinetic energy

$$T = \frac{1}{2}\dot{x}^2$$

and potential energy

$$U(x) = - \int_0^x f(h)dh$$

Note I am following a convention often used in celestial mechanics where kinetic and potential are given per unit mass. In this setting forces are proportional to mass, and the mass drops out of both sides of Newton's equation  $\mathbf{F} = m\mathbf{a}$ . When the dynamics is independent of mass, it can be neglected in the definition of energy!

If the acceleration is downward (toward negative  $x$ ) and constant, the function  $f = -g$ . The potential energy  $U(x) = gx$  and the sign is such that the potential energy is positive for a particle with higher  $x$ . This makes sense if  $g$  is gravitational acceleration, in which case potential energy is larger higher at a higher position.

In one dimension a function  $U(x)$  can be found with  $U' = -f(x)$ . In more than one dimension, forces are vectors but not all vector fields are gradients of a function. A conservative force is one where the force is the gradient of a potential energy function  $\mathbf{F} = -\nabla U$ . Not every vector force is *conservative*. Forces that are conservative are a special case.

### 2.1 Energy is conserved

Energy is the sum of the kinetic and potential energies,  $E = T + U$  with kinetic energy  $T = \dot{x}^2/2$ . Taking the time derivative

$$\frac{d}{dt}(T + U) = \dot{x}\ddot{x} + \frac{dU}{dx}\dot{x} = \dot{x}(\ddot{x} - f(x)) = 0$$

The energy  $E = T + U$  is conserved.

The above conservation law can be generalized for a conservative vector force with  $\nabla U = -\mathbf{F}$

$$\frac{d}{dt}(T + U) = \dot{\mathbf{x}} \cdot \ddot{\mathbf{x}} + \nabla U \cdot \dot{\mathbf{x}} = \dot{\mathbf{x}} \cdot (\ddot{\mathbf{x}} - \mathbf{F}) = 0$$

In more than one dimension, energy is conserved if the force is conservative. We have assumed that force  $\mathbf{F}$  and potential energy  $U$  are independent of time.



In the Hamiltonian viewpoint, conservation of energy follows directly from Hamilton's equations. Taking  $H(q, p)$  but independent of time,

$$\begin{aligned}\frac{dH(q, p)}{dt} &= \frac{\partial H}{\partial q} \dot{q} + \frac{\partial H}{\partial p} \dot{p} \\ &= \dot{p}\dot{q} - \dot{q}\dot{p} = 0.\end{aligned}\tag{6}$$

## 2.2 Phase flow

Energy can be written

$$E(x, \dot{x}) = \frac{1}{2}\dot{x}^2 + U(x)$$

We can consider plotting the motion as a trajectory on a plane defined by  $(x, y)$  where

$$y = \dot{x}.$$

$$E(x, y) = \frac{1}{2}y^2 + U(x)$$

Because energy is conserved, orbits on this plane correspond to level curves of  $E$ . The equations of motion are now

$$\dot{x} = y \quad \dot{y} = f(x) = -\frac{dU}{dx}.\tag{7}$$

A phase curve consists of functions of time such that

$$x = \phi_x(t) \quad y = \phi_y(t)$$

describing a trajectory. These are functions that depend on the initial conditions so we can write  $\mathbf{x} = \phi(t, \mathbf{x}_0)$  where  $\mathbf{x} = (x, y)$  and the initial condition is  $\mathbf{x}_0$ . The equations of motion determine these functions. The trajectory gives a vector field on the phase plane. Level curves of constant energy  $E(x, y)$  are trajectories. We can also refer to trajectories as *orbits*.

### 2.2.1 Example: The Harmonic oscillator

Working with a dimensionless system (spring constant and mass are 1) the force

$$f(x) = -x\tag{8}$$

and energy for a harmonic oscillator

$$E(x, y) = \frac{1}{2}(x^2 + y^2)\tag{9}$$

with  $y = \dot{x}$ . The phase curves for the harmonic oscillator are shown in Figure 2. There is one fixed point, at the origin. The period of oscillations is independent of the amplitude. Orbits are circles.

### 2.2.2 Trajectories can go in either direction

The orbits can circulate about the origin in either direction. This is related to the fact that flipping the direction of time is equivalent to changing the direction of motion (or flipping the sign of the momentum). Let's look at the equations of motion again:

$$\dot{x} = y \quad \dot{y} = -\frac{dU}{dx}. \quad (10)$$

What happens if let  $\tau = -t$ . We find that  $\frac{dx}{d\tau} = -\frac{dx}{dt}$ . But if we also change the sign of  $y$ , then the equations of motion are unchanged. Suppose we have a trajectory  $x(t)$  from  $x = a$  to  $x = b$ . We can flip the arrow of time, to find a trajectory that goes from  $x = b$  to  $x = a$ .

### 2.2.3 What is a fixed point?

For a dynamical system  $\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x})$ . A fixed point is a point that satisfies  $\dot{\mathbf{x}} = \mathbf{0}$ .

The harmonic oscillator has a single fixed point at the origin.

What happens if you move a small distance away from the fixed point? The orbit stays near the fixed point and never gets too far away from the fixed point. The fixed point is **stable** because nearby trajectories do not diverge away from it.

In a Hamiltonian system  $\dot{\mathbf{x}} = \mathbf{0}$  consists of more than one equation. For  $H(q, p)$  the fixed point satisfies  $\dot{q} = 0$  and  $\dot{p} = 0$ . At the fixed point, Hamilton's equations give  $\frac{\partial H}{\partial q} = 0$  and  $\frac{\partial H}{\partial p} = 0$ .

### 2.2.4 Equivalent Hamiltonian model

In 1 dimension with potential  $U(q)$  we can construct a Hamiltonian

$$H(p, q) = \frac{p^2}{2} + U(q)$$

Hamilton's equations give

$$\dot{q} = \frac{\partial H}{\partial p} = p \quad \dot{p} = -\frac{\partial H}{\partial q} = -U'(q) = f(q) \quad (11)$$

The first of these gives  $\ddot{q} = \dot{p}$  which if we combine with the second gives

$$\ddot{q} = f(q)$$

which is consistent with Newton's equations with force per unit mass  $f(q)$ .

For the harmonic oscillator  $f(x) = -x$  and  $U(x) = x^2/2$  giving Hamiltonian

$$H(x, p) = \frac{p^2}{2} + \frac{x^2}{2}.$$




Figure 2: Level curves and phase flow of the Harmonic oscillator.




Figure 3: Level curves and phase flow of the Pendulum.




Figure 4: The difference between librating and oscillating orbits for the pendulum (equation 13). Here the vertical axis is  $y$  and the  $x$  axis corresponds to an angle around each cylinder.

### 2.2.5 Example: Pendulum

Here  $x$  is an angle and the force depends on the sin of the angle.

$$f(x) = -\sin x \quad (12)$$

The energy is

$$E(x, y) = -\cos x + \frac{y^2}{2} \quad (13)$$

with  $\dot{x} = y$ . Whereas  $y$  can take any value, the system is periodic with respect to  $x$ , so we can think of  $x$  as an angle. The dynamical system on the plane can be written

$$\begin{aligned} \dot{x} &= y \\ \dot{y} &= -\sin x \end{aligned}$$

At each time the vector  $\dot{x}, \dot{y}$  gives a *tangent* velocity vector. The level curves for this system are shown in Figure 3. Near the origin the phase curves resemble that of the harmonic oscillator. In fact we can expand the cosine for small  $x$  finding an energy

$$E(x, y) \sim -1 + \frac{x^2}{2} + \frac{y^2}{2} \quad (14)$$

which is a harmonic oscillator. However trajectories approach and leave  $(x, y) = (\pi, 0)$ . This is an **unstable fixed** point. The orbit containing the unstable fixed point is known as the **separatrix** and separates **librating** from oscillating trajectories. Librating trajectories correspond to small oscillations about the origin. Oscillating trajectories corresponds to those at large  $\dot{x}$  (see Figure 4). A pendulum going around and around in a big circle has large  $\dot{x}$  with orbit on the top or bottom of Figure 3. A pendulum has three types of orbits and two types of fixed points (unstable and stable). The types of orbits are the librating orbits (ellipses), **hyperbolic orbits** connecting unstable fixed points that have infinite period (the separatrix), and circulating orbits.

The Hamiltonian for the pendulum consistent with that discussed above is

$$H(\phi, p) = \frac{p^2}{2} - \cos \phi.$$

The potential function  $V(\phi) = -\cos \phi$  gives force  $-\frac{\partial V(\phi)}{\partial \phi} = \sin \phi$ . The equations of motion from Hamilton's equation are

$$\begin{aligned} \dot{\phi} &= p \\ \dot{p} &= -\sin \phi = -\ddot{\phi} \end{aligned}$$

### 2.2.6 Example: Cubic Potential

For a Hamiltonian

$$H(p, q) = \frac{1}{2}p^2 + U(q)$$

fixed points (those with  $\dot{p} = \dot{q} = 0$ ) occur where  $p = 0$  and  $dU/dq = 0$ . If  $U(q)$  is a cubic polynomial (with a non-zero  $q^3$  coefficient) then fixed points are solutions of a quadratic equation in  $q$ . This means there are either zero, one or two real roots. If the potential is continuously rising or falling then there are no fixed points. If there is an inflection point and a single root then the fixed point is unstable. If there are two real roots then one fixed point is unstable and the other stable (see Figure 5). We can draw a bifurcation diagram (see Figure 6). We use a cubic potential

$$U(q) = -aq + q^3/3$$

that is a function only of the parameter  $a$ , then fixed points occur where

$$U'(q) = -a + q^2 = 0.$$

We have two real roots for  $a > 0$  and no real roots for  $a < 0$ . The roots are at

$$q_{fix} = \pm\sqrt{a} \quad \text{for} \quad a > 0$$

and so their separation increases with  $a$ . The bifurcation diagram can be drawn as a function of  $a$ .




Figure 5: A cubic potential shown on the left. Phase flow and level curves shown on the right.




Figure 6: An example of a bifurcation diagram for fixed points of a cubic potential. For a cubic potential  $U(q) = -aq + q^3/3$ , two fixed points exist at  $\pm\sqrt{a}$  when  $a > 0$ . The solid line corresponds to the stable fixed point and the dotted one to the unstable fixed point.

For positive  $a$ , we can expand the Hamiltonian around one of the fixed points. Near  $q = \sqrt{a}$

$$x = q - \sqrt{a}, \quad (15)$$

giving

$$\begin{aligned} H(p, x) &= \frac{p^2}{2} - a(x + \sqrt{a}) + \frac{1}{3}(x + \sqrt{a})^3 \\ &= \frac{p^2}{2} - a(x + \sqrt{a}) + \frac{x^3}{3} + \frac{a^{3/2}}{3} + x^2\sqrt{a} + xa \\ &\approx \frac{p^2}{2} + \sqrt{a}x^2 + \text{constant} \end{aligned} \quad (16)$$

and we have assumed that  $x$  is small. The system looks like a harmonic oscillator. Near  $q = -\sqrt{a}$  and using

$$x = q + \sqrt{a} \quad (17)$$

giving

$$H(p, x) \approx \frac{p^2}{2} - \sqrt{a}x^2 + \text{constant}.$$

Here the level curves are hyperbolic.

Note: Equation 15, 17 with an unchanged  $p$  are canonical transformations and Hamilton's equations are valid in the new coordinate system.

### 2.2.7 Example: The radial degree of freedom in Kepler's problem

We consider the problem of a massless particle in orbit around a massive body of mass  $M$ . We restrict motion to a single plane, the  $x, y$  plane. The potential energy (per unit mass) is a function of radius alone and is  $U = -GM/r$  for mass  $M$  with  $G$  the gravitational constant. Hereafter we drop the phrase *per unit mass* when discussing energy or angular momentum. The kinetic energy

$$T = \frac{1}{2}(\dot{x}^2 + \dot{y}^2) = \frac{1}{2}(\dot{r}^2 + r^2\dot{\phi}^2),$$

where we also use polar coordinates  $r, \phi$ . Neither potential energy or kinetic energy depend on  $\phi$ . That means that the Hamiltonian in polar coordinates does not depend on one of the coordinates,  $\phi$ . Consequently there is a conserved quantity, the angular momentum,  $L = r^2\dot{\phi}$ .

The tangential velocity component of the kinetic energy

$$\frac{1}{2}r^2\dot{\phi}^2 = \frac{L^2}{2r^2}.$$

The radial motion can be described with an effective potential

$$U(r) = \frac{L^2}{2r^2} - \frac{GM}{r} \quad (18)$$

and equations of motion

$$\ddot{r} = -\frac{\partial U}{\partial r}$$

with energy

$$E(\dot{r}, r) = \frac{\dot{r}^2}{2} + \frac{L^2}{2r^2} - \frac{GM}{r}.$$

This system resembles a harmonic oscillator in terms of its level curves. Interestingly we know that the period of radial oscillation in this system only depends on energy  $E$  and is independent of  $L$ . However it is not obvious from the shape of the potential that this is true.






Figure 7: The effective potential for a Keplerian system  $U(r) = \frac{L^2}{2r^2} - \frac{GM}{r}$  drawn for two different values of  $L$ . Circular orbits are located at the potential minimum. Orbital period depends only on energy not angular momentum, though it is not obvious from the shape of the effective potential.




Figure 8: Integrating the area of an orbit. The area enclosed by an orbit with energy  $E$  is  $S(E) = 2 \int_{x_{min}}^{x_{max}} \dot{x} dx$ . There is symmetry about  $\dot{x} = 0$  for an orbit with  $\dot{x} = \pm \sqrt{2(E - U(x))}$ .

## 2.3 Period as a function of area in phase space

Taking a particle in a potential

$$E = \frac{\dot{x}^2}{2} + U(x)$$

We can solve for  $\dot{x}$

$$\dot{x} = \frac{dx}{dt} = \pm \sqrt{2(E - U(x))}$$

giving (for the positive root)

$$dt = \frac{dx}{\sqrt{2(E - U(x))}} \quad (19)$$

and the time between two points  $x_1, x_2$  at  $t_1, t_2$

$$t_2 - t_1 = \int_{x_1}^{x_2} \frac{dx}{\sqrt{2(E - U(x))}} \quad (20)$$

For a periodic orbit (one that comes back to itself) we can compute the area of the orbit  $S(E)$  as a function of energy.

$$S(E) = 2 \int_{x_{min}}^{x_{max}} \dot{x} dx = 2 \int_{x_{min}}^{x_{max}} \sqrt{2(E - U(x))} dx$$

where by symmetry the area for positive  $\dot{x}$  is the same as that for negative  $\dot{x}$ . See Figure 8. If we take the derivative of this with respect to energy

$$\begin{aligned} \frac{\partial S(E)}{\partial E} &= 2 \int_{x_{min}}^{x_{max}} \frac{dx}{\sqrt{2(E - U(x))}} \\ &= 2[t(x_{max}) - t(x_{min})] = P \end{aligned}$$

where  $P$  is the orbital period. In the second step we have used equation 20. So the orbital period

$$P(E) = \frac{\partial S(E)}{\partial E}. \quad (21)$$

Again look at the Kepler system to reflect that it is not obvious that  $P$  is independent of  $L$ .

Multiplying a Hamiltonian by a constant is equivalent to rescaling the Hamiltonian by time. This gives an analogy for why a gradient in energy is related to period.

### 2.3.1 Harmonic and an-harmonic oscillator examples

Consider the harmonic oscillator

$$H(p, q) = \frac{1}{2}(p^2 + q^2).$$

The level curves are circles. In action angle variables the Hamiltonian  $H(I, \theta) = I$  with action variable

$$I = \frac{1}{2}(p^2 + q^2) = \frac{r^2}{2}$$

where  $r$  is the radius on the  $p, q$  plane. The area in phase space

$$S(E) = \pi r^2 = 2\pi I = 2\pi E.$$

The period is  $2\pi$  and consistent with  $S'(E)$ .

Now consider an anharmonic oscillator

$$H(I, \theta) = I + aI^2. \quad (22)$$

The level curves are still circles. However the period for the anharmonic oscillator is dependent on  $I$ . For the anharmonic oscillator we use the quadratic equation with equation 22

$$I = \frac{1}{2a} \left( -1 + \sqrt{1 + 4Ea} \right)$$

and the area as a function of energy is

$$S(E) = 2\pi I = \frac{2\pi}{2a} \left( -1 + \sqrt{1 + 4Ea} \right).$$

We compute

$$P = \frac{\partial S(E)}{\partial E} = 2\pi(1 + 4Ea)^{-\frac{1}{2}} = \frac{2\pi}{1 + 2Ia}.$$

With  $a = 0$  the period is insensitive to the energy and is equal to  $2\pi$  as we expect. We can check the period using Hamiltonian's equation for  $H$  in 22

$$\dot{\theta} = \frac{\partial H(I, \theta)}{\partial I} = 1 + 2aI$$

and a period of

$$P = \frac{2\pi}{\dot{\theta}} = \frac{2\pi}{1 + 2aI}$$

which is consistent with that computed from  $S'(E)$ .

### 2.3.2 Hamiltonians that give the same orbital trajectories

Take a time independent Hamiltonian  $H(\mathbf{p}, \mathbf{q})$  and a different Hamiltonian  $K(\mathbf{p}, \mathbf{q}) = K(H(\mathbf{p}, \mathbf{q}))$ . The level curves for the two Hamiltonians are the same curves. A particle trajectory with energy  $E$  derived with  $H$  is the same trajectory (through phase space) as that with energy  $K(E)$  and with equations of motion derived from  $K$ . However these two orbits do not have the same orbital period.

## 2.4 The separatrix of a pendulum

The time it takes to get from one unstable fixed point to another one is infinite. This can be argued from uniqueness of solutions to our second order differential equation. Also we can compute the energy at the separatrix and show that it is zero at  $x = 0$ . Because phase space is separated by the separatrix, the derivative of the area with respect to energy (equation 21) diverges.

Particles located near a stable fixed point, undergo small oscillations about the fixed point. Particles located near the unstable fixed point exponentially diverge from it or exponentially slowly approach it. For example with

$$H(p, \phi) = \frac{p^2}{2} - \epsilon \cos \phi$$

we can expand near  $\phi = \pi$  with  $\phi = \pi + x$ , giving

$$x \equiv \phi - \pi.$$

For small  $x$

$$\cos(\pi + x) \sim -1 + \frac{x^2}{2}.$$

Hamilton's equations give

$$\begin{aligned} \dot{\phi} &= \dot{x} = \frac{\partial H}{\partial p} = p \\ \dot{p} &= -\frac{\partial H}{\partial \phi} = -\frac{\partial H}{\partial x} = \epsilon x. \end{aligned}$$

Taking the time derivative of  $\dot{x}$  and combining equations we find

$$\ddot{x} = \epsilon x$$

with solution

$$x \propto e^{\pm\sqrt{\epsilon}t}$$

near the unstable fixed point.

The timescale for oscillation about a *stable* fixed point is the oscillation period. The same timescale gives an estimate for how fast an orbit moves away from an unstable or *hyperbolic* fixed point.

## 2.5 On Hamiltonian systems with one degree of freedom

The phase space is two dimensional. For a time independent Hamiltonian, there is a single conserved quantity, the energy or Hamiltonian. All orbits fill at most a 1-dimensional

space. A single quantity (energy) defines a trajectory. The system is said to be *integrable*. All smooth forces are *conservative*.

When we consider two degrees of freedom, many of these statements are no longer true. Not all forces are conservative. Phase space is 4 dimensional and so one conserved quantity is not enough to define a trajectory. An interesting bridge between the one-dimensional and two dimensional systems are periodically forced systems such as the periodically forced pendulum or the kicked rotor that is the physical analog for the standard map.

### 3 Relation between Lagrangian and Hamiltonian Descriptions

#### 3.1 Extremum of a path

A Lagrangian is a function  $\mathcal{L}(q, \dot{q}, t)$ . We consider a path  $\gamma$  that gives  $q(t)$  for  $t_1 \leq t \leq t_2$ . The equations of motion are found by minimizing the integral along the path

$$\Phi(\gamma) = \int_{t_1}^{t_2} \mathcal{L}(q, \dot{q}, t) dt$$

Here  $q(t)$ ,  $\dot{q}(t)$ . By using calculus of variations we can show that Lagrange's equation is true on the extremum curve.

We differentiate this function by changing the path by an amount  $q(t) \rightarrow q(t) + h(t)$  (see Figure 9). We chose  $q_1(t_1)$  and  $q_2(t_2)$  and assert that the ends of the path remain fixed;  $h(t_1) = h(t_2) = 0$ .

$$\begin{aligned} \Phi(\gamma + h) - \Phi(\gamma) &= \int_{t_1}^{t_2} \left[ \mathcal{L}(q + h, \dot{q} + \dot{h}, t) - \mathcal{L}(q, \dot{q}, t) \right] dt \\ &= \int_{t_1}^{t_2} \left[ \frac{\partial \mathcal{L}}{\partial q} h + \frac{\partial \mathcal{L}}{\partial \dot{q}} \dot{h} \right] dt + O(h^2) \end{aligned} \quad (23)$$

Integrating by parts

$$\int_{t_1}^{t_2} \frac{\partial \mathcal{L}}{\partial \dot{q}} \dot{h} dt = - \int_{t_1}^{t_2} \frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{q}} h dt + \left( h \frac{\partial \mathcal{L}}{\partial \dot{q}} \right) \Big|_{t_1}^{t_2}$$

We insert this back into equation 23

$$\Phi(\gamma + h) - \Phi(\gamma) = \int_{t_1}^{t_2} \left[ \frac{\partial \mathcal{L}}{\partial q} - \frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{q}} \right] h dt + \left( h \frac{\partial \mathcal{L}}{\partial \dot{q}} \right) \Big|_{t_1}^{t_2}$$

Because we have asserted that the ends of the path remain fixed,  $h = 0$  on the ends of the path, and the rightmost term must be zero. Hence the extremum of the integral  $\Phi(\gamma)$  is a

path on which Lagrange's equations are satisfied or

$$\frac{\partial \mathcal{L}}{\partial q} - \frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{q}} = 0$$




Figure 9: Paths between points  $q_1(t_1)$  and  $q_2(t_2)$

### 3.2 Lagrangian's equations for a Newtonian system

For a mechanical system  $L = T(\dot{q}) - U(q)$  and Lagrange's equations are equivalent to Newton's equations.

$$L(q, \dot{q}, t) = \frac{1}{2} \dot{q}^2 - U(q)$$

$$\frac{\partial L}{\partial \dot{q}} = \dot{q} \tag{24}$$

$$\frac{\partial L}{\partial q} = -\frac{dU}{dq} = f(q) \tag{25}$$

Hence Lagrange's equations give

$$\ddot{q} = f(q)$$

which is Newton's equation.

### 3.3 Hamilton's equations for a Newtonian system

The Hamiltonian for a Newtonian system is

$$H(p, q) = \frac{p^2}{2} + U(q) \tag{26}$$




Figure 10: A figure that helps visualize the Legendre transform. The transform  $g(p)$  of a convex function,  $f(x)$  is  $g(p) = \sup_x (xp - f(x))$ . In this figure we choose a  $p$ . The blue line shows the line  $y = px$ . This matches the slope of  $f(x)$  at a point  $x'$ . The function  $F(x, p) = xp - f(x)$  is maximized at  $x'$  where  $f'(x') = p$ . The thick blue dashed line is  $y = b + px$  with intercept  $b = -(px' - f(x'))$ . The transform takes  $(x, f(x))$  and gives (slope, intercept(slope)) or  $(p, g(p))$  using lines tangent to  $f$ . We define a new function  $g(p)$  that is the negative of the  $y$ -intercept of the tangent line to  $f$  that has slope  $p$ .

Hamilton's equations give

$$\dot{q} = \frac{\partial H}{\partial p} = p$$

$$\dot{p} = -\frac{\partial H}{\partial q} = -U' = f(q)$$

where  $f(q)$  is the force associated with the potential  $U(q)$ . Taking the time derivative of  $\dot{q}$  we find

$$\ddot{q} = f(q)$$

which is Newton's equation.

Both Hamiltonian and Lagrangian viewpoints give Newton's equations. In the next section we will show how a Lagrangian can be turned into a Hamiltonian and how Lagrange's equations are consistent with Hamilton's equations.

### 3.4 Legendre Transforms

A Lagrangian system can be converted into a Hamiltonian system using a Legendre transform.

A Legendre transform transforms functions on a vector space to functions on a dual space. It transforms functions of a coordinate  $x$  to functions that are dependent on the slope of  $f(x)$ . Let  $y = f(x)$  and look at Figure 10. We specify that  $f(x)$  is convex so that  $f''(x) > 0$  everywhere. We chose a value  $p$  for a slope and draw the line  $px$ . We look for the point where  $px - f(x)$  is a maximum. Let us define a function  $F(p, x) = px - f(x)$  that gives this distance. The point where  $F(p, x)$  is a maximum is the point where the vertical distance between  $f(x)$  and  $px$  is the largest. The condition that  $F(p, x)$  is an extremum at  $x'$  means that  $\frac{\partial F}{\partial x}|_{x'} = 0$ . And this implies that  $f'(x') = p$ . The point  $x'$  is special because  $f'(x') = p$  so  $p$  is the slope of  $f(x)$  at  $x'$ . On the figure the special point is labelled as  $x'$ . For any  $p$  this point,  $x'$  is unique as  $f(x)$  is convex. We define a new function  $g(p) = F(p, x') = px' - f(x')$ . This function gives the negative of the  $y$  intercept of the tangent line to  $f$  that has slope  $p$ . The Legendre transform is  $f(x) \rightarrow g(p)$ . Starting with  $(x, f(x))$  the Legendre transform gives a new function  $(p, g(p))$  where  $p$  is a slope and  $g(p)$  is the negative of the  $y$  intercept of the tangent to  $f$  that has slope  $p$ .

$$(x, f(x)) \rightarrow (p, g(p))$$

$$(\text{position, function}) \rightarrow (\text{slope, } -\text{intercept})$$

The Legendre transform can also be written

$$g(p) = \sup_x (xp - f(x)). \quad (27)$$

The Legendre transform of the Legendre transform gives back the original function (except for a constant). Figure 11 shows this with the points on the  $y$  axis representing the legendre



transform. The function  $G(p, x) = px - g(p)$  is maximized where the lines are tangent to  $f(x)$ .

$$f(x) = \sup_p (px - g(p)). \quad (28)$$




Figure 11: A figure that helps visualize the inverse Legendre transform. The y-intercept is  $-g(p)$ . The function  $G(p, x) = px - g(p)$  is an extremum at a slope  $p'$  where  $G(p', x) = f(x)$ . We can write  $f(x) = \sup_p [px - g(p)]$ .

### 3.4.1 Example: Legendre transform of sqrt(x)

Let's take the Legendre transform of  $f(x) = x^{1/2}$ . Construct

$$F(x, p) = px - f(x) = px - \sqrt{x}$$

We look for the extremum of this function w.r.t.  $x$

$$\frac{\partial F(x, p)}{\partial x} = p - f'(x) = p - \frac{1}{2}x^{-1/2}$$

Set this to zero. The extremum is at

$$p(x') = \frac{1}{2\sqrt{x'}} \quad \text{or} \quad x'(p) = \frac{1}{4p^2}$$

Insert this back into  $F$

$$g(p) = F(x'(p), p) = \frac{p}{4p^2} - \frac{1}{2p} = -\frac{1}{4p}$$

So the Legendre transform is

$$g(p) = -\frac{1}{4p}$$

### 3.5 Constructing a Hamiltonian from a Lagrangian

Recall our Lagrangian is a function of  $\dot{q}, q, t$  or  $\mathcal{L}(q, \dot{q}, t)$ . Let's for the moment ignore the  $q, t$  degrees of freedom so  $\mathcal{L}(\dot{q})$ . Instead of using  $\dot{q}$  as a variable let us use slope  $p = \frac{\partial \mathcal{L}(\dot{q})}{\partial \dot{q}}$  as our variable. The Legendre transform gives us the negative of the y intercept of the tangent line with slope  $p$ . This is

$$H(p) = \sup_{\dot{q}'} [p\dot{q}' - \mathcal{L}(\dot{q}')] \quad (29)$$

with  $\dot{q}'$  the special value of  $\dot{q}$  where  $p = \frac{\partial \mathcal{L}(\dot{q})}{\partial \dot{q}}$ . The special  $\dot{q}'$  is a function of  $p$ .

Now recalling that  $\mathcal{L}$  was also a function of  $q$ , we can write

$$H(p, q, t) = p\dot{q}(p) - \mathcal{L}(q, \dot{q}(p), t) \quad (30)$$

with the condition for  $\dot{q}(p)$

$$p = \frac{\partial \mathcal{L}(q, \dot{q}(p), t)}{\partial \dot{q}}. \quad (31)$$

Let's take the partial derivative of equation 30

$$\frac{\partial H}{\partial p} = \dot{q}(p) + p \frac{d\dot{q}(p)}{dp} - \frac{\partial \mathcal{L}}{\partial \dot{q}} \frac{d\dot{q}(p)}{dp}$$

The right two terms cancel (because of equation 31) giving

$$\frac{\partial H}{\partial p} = \dot{q}$$

one of Hamilton's equations.

To find the second of Hamilton's equations we need to use Lagrange's equation

$$\frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{q}} = \frac{\partial \mathcal{L}}{\partial q}.$$

Our definition of  $H$  (in equation 29) implies that  $\frac{\partial H}{\partial q} = -\frac{\partial \mathcal{L}}{\partial q}$ , hence Lagrange's equation gives

$$\dot{p} = -\frac{\partial H}{\partial q}$$

and we have recovered both of Hamilton's equations.

### 3.5.1 Example: Hamiltonian for a Newtonian system in one dimension

For a Lagrangian

$$\mathcal{L}(q, \dot{q}, t) = \frac{1}{2}\dot{q}^2 - U(q)$$

The Legendre transformation is with respect to the variable  $\dot{q}$ . For the moment we can ignore  $q$ . We consider the function

$$F(\dot{q}, p) = p\dot{q} - \mathcal{L}(\dot{q}) = p\dot{q} - \frac{1}{2}\dot{q}^2 + U$$

The extremum of this function

$$\frac{\partial F}{\partial \dot{q}} = p - \dot{q} = 0$$

occurs for a momentum

$$p = \frac{\partial L}{\partial \dot{q}} = \dot{q}$$

We then insert this particular value of  $\dot{q}(p)$  back into the function  $F$  (which we now call  $H$ )

$$H(p, q) = p\dot{q}(p) - \mathcal{L}(q, \dot{q}(p), t) = \frac{1}{2}p^2 + U(q)$$

## 3.6 Geometrical view of a Lagrangian path integral

Consider the integral giving distance along a path that is between points  $q_1, q_2$

$$\int_{q_1}^{q_2} ds$$

where  $ds$  gives an infinitesimal path length. For example path lengths in cylindrical coordinates

$$ds^2 = dr^2 + r^2 d\theta^2 + dz^2$$

giving metric elements  $g_{rr} = g_{zz} = 1, g_{\theta\theta} = r^2$  and off-diagonal elements that are zero. On the surface of a sphere of radius  $R$

$$ds^2 = R^2 d\phi^2 + R^2 \sin^2 \phi d\theta^2$$

If we have a metric  $g(q)$  then we can write

$$ds^2 = g_{\mu\nu} dq^\mu dq^\nu$$

where I have used summation notation. We can rewrite this

$$ds = \sqrt{g_{\mu\nu} \frac{dq^\mu}{dt} \frac{dq^\nu}{dt}} dt$$

Integrating

$$\int_{q_1}^{q_2} ds = \int_{t_1}^{t_2} \sqrt{g_{\mu\nu} \frac{dq^\mu}{dt} \frac{dq^\nu}{dt}} dt$$

If we minimize the path length then we also minimize a Lagrangian system with

$$L(q, \dot{q}, t) = \sqrt{g_{\mu\nu} \dot{q}^\mu \dot{q}^\nu} \quad (32)$$

Geodesics are then the same as dynamical paths. If we start with a metric we can find a Lagrangian system. The inverse is not so easy. Given a Lagrangian system, it may not be straightforward to write the problem in terms of a metric.

Applying Lagrange's equation to 32, one can derive geodesic equations. When re-written using Christoffel systems (that depend on derivatives of the metric), we obtain the geodesic equation of general relativity.

### 3.7 Pathological Legendre Transformations

Can it be impossible to make the transformation from a Lagrangian system to a Hamiltonian system? The Legendre transformation of a function can only be done uniquely if the function is convex. A twice differentiable function is convex if its second derivative is always positive.

For example consider a non-convex function that has the same slope at two different points. In this case  $p = f'(x)$  is not uniquely inverted! There would be two  $x$  values giving the same slope. Another example is  $H(I, \theta) = I\omega$ . This is a line and so not convex. Oddly I find that  $H(I, \theta) = I + aI^2$  is also badly behaved and gives a zero Lagrangian everywhere. So one can find coordinate systems in which the transformation between Hamiltonian and Lagrangian prescriptions is not possible.

Are there examples of Lagrangians and Hamiltonians for physics systems that are not convex? Al Shapere's (and Wilczek's) time crystals are an entertaining example of this, see *Classical Time Crystals*, Shapere, A. Wilczek F. 2012, Phys. Rev. Lett. 109, 160402

Consider

$$\mathcal{L}(q, \dot{q}) = \frac{\dot{q}^4}{4} - \frac{\dot{q}^2}{2} - V(q) \quad (33)$$

(see Figure 12) To create a Hamiltonian we need momentum that satisfies

$$p(q_*) = \frac{\partial \mathcal{L}(\dot{q}_*)}{\partial \dot{q}}$$

Inverting this  $q_*(p)$ , giving us a Hamiltonian

$$H(p, q) = p\dot{q}_*(p) - \mathcal{L}(q, \dot{q}_*(p))$$

For the Lagrangian in equation 33

$$\frac{\partial \mathcal{L}}{\partial \dot{q}} = \dot{q}^3 - \dot{q} \quad (34)$$

Giving

$$p = \dot{q}_*^3 - \dot{q}_* \quad (35)$$

It may be useful to compute

$$\frac{\partial^2 \mathcal{L}}{\partial \dot{q}^2} = 3\dot{q}^2 - 1$$

Equation 35 is not uniquely invertible for  $q \in [-1/\sqrt{3}, 1/\sqrt{3}]$ .

The Hamiltonian written in terms of  $\dot{q}_*$  is

$$\begin{aligned} H(p, q) &= p\dot{q}(p) - \mathcal{L}(\dot{q}(p), q, t) \\ &= (\dot{q}_*^3 - \dot{q}_*)\dot{q}_* - \frac{\dot{q}_*^4}{4} + \frac{\dot{q}_*^2}{2} + V(q) \\ &= \frac{3}{4}\dot{q}_*^4 - \frac{\dot{q}_*^2}{2} + V(q). \end{aligned}$$

The Hamiltonian (energy) is minimized at  $\dot{q}_* = 1/\sqrt{3}$  or at  $\dot{q}_* = 0$ . But the non-zero value is precisely that value where the second derivative of the Lagrangian is zero (and so the function not convex) and a position where  $p$  is nearly independent of  $\dot{q}$  and so  $p(q_*)$  not invertible.

Let us look at the equations of motion derived from Lagrange's equation:

$$\begin{aligned} \frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{q}} &= \frac{\partial \mathcal{L}}{\partial q} \\ \ddot{q} \frac{\partial^2 \mathcal{L}}{\partial \dot{q}^2} &= \frac{\partial V}{\partial q} \\ \ddot{q}(3\dot{q}^2 - 1) &= \frac{\partial V}{\partial q} \end{aligned}$$

We find that we cannot solve for  $\ddot{q}$  at the same location that minimizes the energy. So breakdown of the equations of motion occurs at the same location as time reversal symmetry is broken (and the same location as our Legendre transformation fails). Life gets more interesting in the quantum setting, and perhaps there are ways around these problems making it possible to break time-reversal symmetry in the ground state.




Figure 12: A non convex Lagrangian. The momentum  $p$  is not a unique function of  $\dot{q}$  giving a multivalued Hamiltonian.

### 3.8 Phase Space Action

Consider a path in phase space  $p(t), q(t)$  and the action

$$S(p, q) = \int (p\dot{q} - H(p, q))dt.$$

Variation of  $S$  with respect to  $q$  gives

$$\begin{aligned} \frac{d}{dt} \frac{\partial}{\partial \dot{q}} (p\dot{q} - H) &= \frac{\partial}{\partial q} (p\dot{q} - H) \\ \frac{d}{dt} p &= -\frac{\partial H}{\partial q} \end{aligned}$$

Variation of  $S$  with respect to  $p$  gives

$$\begin{aligned} \frac{d}{dt} \frac{\partial}{\partial \dot{p}} (p\dot{q} - H) &= \frac{\partial}{\partial p} (p\dot{q} - H) \\ 0 &= \dot{q} - \frac{\partial H}{\partial p} \end{aligned}$$

giving Hamilton's equations. The phase-space action yields Hamilton's equations as the extremal condition.

## 4 Stability of fixed points

The Hamiltonian is a recipe for computing trajectories. We search for points that don't move. These would have  $\dot{p} = \dot{q} = 0$  and we can call them *fixed points* and if they are stable fixed points then we refer to them as *equilibrium points*. For a Hamiltonian system in one degree of freedom (2d phase space), fixed points have

$$\dot{p} = 0 \quad \dot{q} = 0$$

The location  $(p, q)$  of fixed points can be found by solving Hamilton's equations equal to zero

$$\frac{\partial H(p, q)}{\partial p} = \frac{\partial H(p, q)}{\partial q} = 0$$

Suppose we have  $(p_0, q_0)$  that is a fixed point. We can consider the equations of motion near this point. What are the trajectories like near the fixed point. If the trajectories whisk a particle far away then we can refer to the fixed point as *unstable*. If trajectories remain nearby for all time then the fixed point is *stable*. The fixed point is also an *equilibrium* point. Consider the symmetric matrix

$$\mathbf{M} = \begin{pmatrix} \frac{\partial^2 H}{\partial q^2} & \frac{\partial^2 H}{\partial p \partial q} \\ \frac{\partial^2 H}{\partial p \partial q} & \frac{\partial^2 H}{\partial p^2} \end{pmatrix} \quad (36)$$

evaluated at  $p_0, q_0$ . If we expand the Hamiltonian near the fixed point  $q = q_0 + x$ ,  $p = p_0 + y$

$$H(x, y) = H(p_0, q_0) + \frac{\partial^2 H(p_0, q_0)}{\partial q^2} \frac{x^2}{2} + \frac{\partial^2 H(p_0, q_0)}{\partial p^2} \frac{y^2}{2} + \frac{\partial^2 H(p_0, q_0)}{\partial p \partial q} xy \quad (37)$$

or

$$H(x, y) = \frac{1}{2} \begin{pmatrix} x & y \end{pmatrix} \begin{pmatrix} M_{xx} & M_{xy} \\ M_{xy} & M_{yy} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$$

where we have dropped the constant term. Our Hamiltonian system can also be written

$$H(\mathbf{x}) = \frac{1}{2} \mathbf{x}^t \mathbf{M} \mathbf{x} \quad (38)$$

where we have defined a vector  $\mathbf{x} = (x, y)$  and  $\mathbf{x}^t$  is the transpose of  $\mathbf{x}$ . Here  $M$  is the matrix in equation 36 evaluated at the fixed point. Or in terms of a dot product

$$H(\mathbf{x}) = \frac{1}{2} \langle \mathbf{x} | \mathbf{M} \mathbf{x} \rangle$$

We can also write this in summation notation

$$H(\mathbf{x}) = \frac{1}{2} x_i M_{ij} x_j$$

where repeated indices are summed. Taking the gradient

$$\begin{aligned} \frac{\partial H}{\partial x_k} &= \frac{\partial}{\partial x_k} \sum_{ij} \frac{1}{2} x_i M_{ij} x_j \\ &= \frac{1}{2} \sum_{jk} [\delta_{ik} M_{ij} x_j + \delta_{jk} x_i M_{ij}] \\ &= \frac{1}{2} \left[ \sum_j M_{kj} x_j + \sum_i M_{ik} x_i \right] \\ &= \frac{1}{2} \left[ \sum_j M_{kj} x_j + \sum_i M_{ki} x_i \right] \\ &= \frac{1}{2} \left[ \sum_j M_{kj} x_j + \sum_j M_{kj} x_i \right] \\ &= \sum_j M_{kj} x_j \end{aligned} \quad (39)$$

In vector notation

$$\nabla H = \mathbf{M} \mathbf{x}$$



The matrix is symmetric and that means it has two real eigenvalues. Expanding near the fixed point preserves Hamilton's equations with  $x$  acting as a coordinate and  $y$  acting as a momentum. So the coordinate transformation to  $(x, y)$  is said to be *canonical*.

We can diagonalize the matrix. In a rotated coordinate system we can write the Hamiltonian as

$$H(p, q) = \frac{1}{2} (\lambda_1 p^2 + \lambda_2 q^2) \quad (40)$$

where  $\lambda_1, \lambda_2$  are the eigenvalues. If both eigenvalues are positive or both eigenvalues are negative then the system behaves like a harmonic oscillator and has librations about the fixed point with frequency

$$\omega = \sqrt{\lambda_1 \lambda_2}$$

The fixed point is stable. If the product of the eigenvalues is negative then evolution is exponential with trajectories diverging exponentially along one eigenvector and heading toward the fixed point along the other one. The fixed point is unstable.

**Remark** Taking into account the units in equation 40 There is only one way to make a frequency from the eigenvalues  $\lambda_1, \lambda_2$ . This corresponds to the frequency of oscillation about a stable fixed point. For an unstable fixed point, this frequency corresponds to that for exponential behavior near the fixed point.

**Remark** The harmonic oscillator (figure 2) contains a single stable fixed point at the origin. The pendulum has examples of both types of fixed points (see Figure 3). The unstable fixed point are at  $x = \pm\pi$  and the stable ones at  $x = 0, 2\pi$ .

**Question:** Why do we ignore the possibility that one of the eigenvalues is zero?

Answer: Volume conservation.

Let us take another look at equation 38. Using Hamilton's equations in the form

$$\dot{\mathbf{x}} = \boldsymbol{\omega} \nabla H$$

with

$$\boldsymbol{\omega} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

we find

$$\dot{\mathbf{x}} = \boldsymbol{\omega} \mathbf{M} \mathbf{x}$$

with  $\mathbf{M}$  defined as the Hessian matrix of the Hamiltonian evaluated at the fixed point (equation 36).

$$\ddot{\mathbf{x}} = \boldsymbol{\omega} \mathbf{M} \boldsymbol{\omega} \mathbf{M} \mathbf{x}$$

For the 2d system the matrix

$$(\omega \mathbf{M})^2 = -(\det\{\mathbf{M}\})\mathbf{I}$$

where  $\mathbf{I}$  is the identity matrix. If the determinant is positive (two positive or two negative eigenvalues) then the fixed point is stable and the frequency of oscillation is

$$\nu = \sqrt{\det\{\mathbf{M}\}} = \sqrt{\lambda_1 \lambda_2}$$

If the determinant is negative then solutions are exponentially dependent on time with timescale

$$t_{exp} = \frac{1}{\sqrt{|\det\{\mathbf{M}\}|}}$$

## 4.1 Lyapunov and asymptotic stability

*Stability* can be defined more rigorously in the following way. Suppose  $\mathbf{x}_f$  is our fixed point and  $\phi(t, \mathbf{x}_0)$  a trajectory as a function of time with initial condition  $\mathbf{x}_0$ . A fixed point is stable (Lyapunov stable) if for every  $\epsilon > 0$  there is a  $\delta$  such that

$$\|\mathbf{x}_f - \phi(t, \mathbf{x}_0)\| < \epsilon$$

for all  $t$  and  $\mathbf{x}_0$  that satisfy

$$\|\mathbf{x}_f - \mathbf{x}_0\| < \delta$$

This means that trajectories never get far away from the fixed point ever, either in future or past. An example of trajectories near a stable fixed point is shown in Figure 13.

A fixed point is asymptotically stable if in addition to being Lyapunov stable, all trajectories converge to that point. Hamiltonian systems do not contain asymptotically stable fixed points because volume in phase space is conserved. However, dissipative systems can contain asymptotically stable fixed points. These are often described as *attracting*.

Sometimes it is possible construct a function that is always decreasing. For example, in a dissipative system, the energy would always be decreasing. A function that is always decreasing is called a *Lyapunov function* and can be used to establish boundaries of conditions that decay to a particular fixed point.

## 5 Dimensional rescaling for Hamiltonians

Consider kinetic and potential energy per unit mass. This is appropriate for a low mass particle in orbit about the Sun for example. In this setting we can compute Energy per unit mass as the sum of kinetic and potential energy per unit mass. Our Hamiltonian




Figure 13: A stable fixed point. The level curves of a 1dimensional harmonic oscillator are shown at the bottom. Trajectories are shown with time in the third dimension. The fixed point is shown with a blue trajectory. Nearby orbits always remain in the vicinity of the stable fixed point.

would have units of energy per unit mass or velocity<sup>2</sup>. Since we are ignoring mass we can take momentum as having units of velocity.

$$H = T + V.$$

Hamilton's equations for  $H(p, q, t)$

$$\frac{\partial H}{\partial p} = \dot{q} \quad (41)$$

$$\frac{\partial H}{\partial q} = -\dot{p} \quad (42)$$

For a free particle  $H = \frac{p^2}{2}$ . If I multiply  $H$  by a constant  $H = \frac{ap^2}{2}$  I would find  $\dot{q} = ap$  instead of  $\dot{q} = p$ . But if I replace  $dt$  with  $d\tau/a$  (setting  $t = \tau/a$ ) I find that  $\frac{dq}{dt} = \frac{dq}{d\tau}a$  and I can recover the constant  $a$ . Essentially multiplying the Hamiltonian by a constant  $a$  is equivalent to rescaling time by a factor  $1/a$ . Let's do this more formally. Consider the transformation  $\frac{d\tau}{dt} = a$  which is consistent with  $\tau = ta$ .

$$\dot{q} = \frac{dq}{dt} = \frac{dq}{d\tau} \frac{d\tau}{dt} = a \frac{dq}{d\tau}$$

$$\dot{p} = a \frac{dp}{d\tau}$$

In a new coordinate system but using  $\tau$  as time instead of  $t$  our new Hamiltonian

$$H(p, q, \tau) = H(p, q, t)a.$$

Momentum and coordinates are not changed with this rescaling of time.

A numerical technique known as *regularization* exploits the fact that multiplying a Hamiltonian by time is like rescaling time. Instead of multiply by a constant, a Hamiltonian is multiplied by a function that might depend on coordinates. A new Hamiltonian system is created with different units of time. For example it is desirable to have short timesteps near a large point mass and large timesteps distant from a point mass. Regularized systems can be used to create more accurate integrators for high eccentricity stars near black holes, for example.

Regard Hamilton's equations again. If I replace  $q$  by  $aQ$  and  $p$  by  $P/a$  then Hamilton's equations are unchanged. Rescaling momentum and inversely scaling coordinates by a constant preserves Hamilton's equations which involve derivatives with respect to both  $p, q$ . This is counter intuitive as one would think that rescaling length would affect both  $p, q$  by the same amount. However if we consider phase space volume as being something that is invariant, then it makes more sense.

Let's multiply the Hamiltonian by a more interesting time dependent function

$$H(p, q, t) = e^{-\lambda t} h(q, p)$$

Hamiltonian's equation give

$$\begin{aligned}\frac{dq}{dt} &= e^{-\lambda t} \frac{\partial h}{\partial p} \\ \frac{dp}{dt} &= -e^{-\lambda t} \frac{\partial h}{\partial q}.\end{aligned}$$

Because the Hamiltonian is time dependent, it is not conserved. Let

$$\tau = -e^{-\lambda t},$$

giving  $\frac{d\tau}{dt} = \lambda e^{-\lambda t}$  and

$$\frac{dt}{d\tau} = \lambda e^{\lambda t}.$$

$$\begin{aligned}\frac{dq}{d\tau} &= \frac{dq}{dt} \frac{dt}{d\tau} = \lambda \frac{\partial h}{\partial p} \\ \frac{dp}{d\tau} &= \frac{dp}{dt} \frac{dt}{d\tau} = -\lambda \frac{\partial h}{\partial q}.\end{aligned}$$

We recover the behavior of a time independent system with Hamiltonian  $\lambda h(p, q)$ . This example shows that you can rescale time with a time dependent function. The behavior of a dissipative system can be mimicked (or approximated with) with a time independent Hamiltonian system.

## 6 Liouville's Theorem and Phase wrapping

Hamilton's equations treat coordinates and momenta differently. However we can consider the dynamics of the system with a  $2N$  vector  $\mathbf{x} = (q_1, q_2, \dots, q_n, p_1, p_2, \dots, p_n)$  where  $N$  is the dimension of  $\mathbf{p}$  and  $\mathbf{q}$ . Our Hamiltonian is  $H(\mathbf{x})$ . Hamilton's equations involve partial derivatives of  $H$  or the gradient operator

$$\nabla H = \left( \frac{\partial H}{\partial q_1}, \frac{\partial H}{\partial q_2}, \dots, \frac{\partial H}{\partial q_n}, \frac{\partial H}{\partial p_1}, \frac{\partial H}{\partial p_2}, \dots, \frac{\partial H}{\partial p_n} \right)$$

Using this notation Hamilton's equation can be written as

$$\dot{\mathbf{x}} = \boldsymbol{\omega} \nabla H \tag{43}$$

where the matrix

$$\boldsymbol{\omega} = \begin{pmatrix} 0 & \mathbf{I} \\ -\mathbf{I} & 0 \end{pmatrix} \tag{44}$$

has dimension  $2N \times 2N$ , and  $\mathbf{I}$  is an identity matrix of dimension  $N$ . The matrix  $\omega$  is antisymmetric so  $\omega_{ij} = -\omega_{ji}$ . The equations of motion (or Hamilton's equations) in summation notation are

$$\dot{x}_i = \omega_{ij} H_{,j}$$

where shorthand  $H_{,j} = \frac{\partial H}{\partial x_j}$ .

Consider the divergence of the flow

$$\nabla \cdot \dot{\mathbf{x}} = \partial_i \dot{x}_i = \partial_i \omega_{ij} H_{,j} = \omega_{ij} H_{,ij}$$

where the right hand side I have used summation notation. The antisymmetry of  $\omega$  implies that

$$\nabla \cdot \dot{\mathbf{x}} = 0$$

Let us show this when we have 1 degree of freedom and a 2-dimensional phase space

$$\begin{aligned} \dot{\mathbf{x}} &= (\dot{q}, \dot{p}) = \left( \frac{\partial H}{\partial p}, -\frac{\partial H}{\partial q} \right) \\ \nabla \cdot \dot{\mathbf{x}} &= \frac{\partial \dot{q}}{\partial q} + \frac{\partial \dot{p}}{\partial p} = \frac{\partial^2 H}{\partial p \partial q} - \frac{\partial^2 H}{\partial q \partial p} = 0 \end{aligned}$$

When  $\nabla \cdot \dot{\mathbf{x}} = 0$ , the flow is *incompressible* in phase space, and we have proved Liouville's theorem.

## 6.1 The meaning of incompressible flow

Let us expand on the connection between being divergence free and being incompressible. Consider a distribution of particles in phase space. The number of particles per unit volume would be  $\rho(\mathbf{p}, \mathbf{q})$  or  $\rho(\mathbf{x})$ . This the number of particles at position  $\mathbf{x}$  in a region with volume  $dV = dx^{2N} = dp^N dq^N$ . We can consider the flux  $\rho \mathbf{v}$  of particles out of each surface of a box that has volume  $dV$ . Integrating the flux over all surfaces of a box gives us the number of particles leaving the box per unit time.

$$\frac{dN}{dt} = \int_S \rho \mathbf{v} d\mathbf{A}$$

over the surface,  $S$ , of the box. Here the velocity  $\mathbf{v} = \dot{\mathbf{x}}$ . Using Gaus' law we can rewrite this as

$$\frac{dN}{dt} = \int_V \nabla \cdot (\rho \mathbf{v}) dV$$

If particles are leaving the box then the number density in the box must decrease.

$$\frac{dN}{dt} = - \int \frac{\partial \rho}{\partial t} dV$$

We find a conservation law

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{v}) = 0$$

which can be recognized as conservation of mass for a fluid flow. Consider a bowling ball moving through water. As the bowling ball moves past an observer, it would appear that the density at some location is changing. However the bowling ball is not being compressed. The problem is that the above equation is written respect to a fixed coordinate system and we need to think about the density in a volume element that is moving with the fluid.

We can rewrite this equation as

$$\begin{aligned} \frac{\partial \rho}{\partial t} + (\mathbf{v} \cdot \nabla) \rho &= -\rho \nabla \cdot \mathbf{v} \\ \frac{D\rho}{Dt} &= -\rho \nabla \cdot \mathbf{v} \end{aligned} \tag{45}$$

where I have used

$$\frac{D}{Dt} \equiv \frac{\partial}{\partial t} + \mathbf{v} \cdot \nabla$$

is the advective derivative or that moving with the fluid. Consider  $\rho(\mathbf{x}, t)$  as it varies

$$\begin{aligned} \frac{d\rho}{dt} &= \frac{\partial \rho}{\partial x} \dot{x} + \frac{\partial \rho}{\partial y} \dot{y} + \frac{\partial \rho}{\partial z} \dot{z} + \frac{\partial \rho}{\partial t} \\ &= \mathbf{v} \cdot \nabla \rho + \frac{\partial \rho}{\partial t} \end{aligned}$$

The equation 45 implies that when  $\nabla \cdot \mathbf{v} = 0$  then  $\frac{D\rho}{Dt} = 0$  and the density of a distribution of particles remains fixed even as the distribution of particles is deformed.

## 6.2 Phase wrapping

If we start with a bunch of evenly distributed particles and look at the area covered by then for a 2D phase space (1 degree of freedom Hamiltonian) then the area covered by these particles remains constant.

For a harmonic oscillator the period of a orbit is independent of the amplitude. The phase flow could cause a distribution of particles originally located in a small region of phase space distant from the origin to rotate about the origin. The area and shape of the distribution would remain content but the central value of  $x, \dot{x}$  for the distribution would move along the orbit (see Figure 14). The harmonic oscillator  $H(p, q) = \frac{1}{2}(p^2 + \omega^2 q^2)$  can also be written

$$H(I, \phi) = I\omega$$

which has  $\dot{\phi} = \omega$  independent of amplitude or  $I$ . On the level curve plot  $I$  sets the distance from the origin.

However in many oscillating systems the period of the orbit depends on the amplitude. This is true for the pendulum and even at moderate amplitude for the anharmonic oscillator with a quartic potential  $U(x) = x^2 + ax^4$ . The area enclosed by the distribution remains constant as the flow is divergence-free. A distribution of particles originally in a small region in phase space is sheered out because particles have different amplitudes and so different oscillation frequencies. This kind of system can be described with a Hamiltonian

$$H(I, \phi) = I\omega + aI^2$$

giving  $\dot{\phi} = \omega + aI$ . The period depends on the distance from the origin in Figure 15a or the  $I$  momentum. See Figure 15 for an illustration. This phenomena is called *phase wrapping* because particles with different amplitudes diverge in orbital *phase*.




Figure 14: The distribution of particles at three different times are shown as red regions. On left: the flow of a harmonic oscillator  $H(q, p) = \frac{1}{2}(p^2 + q^2)$ . There is no phase wrapping because the period is independent of amplitude. On right: for an anharmonic oscillator where the period depends on the amplitude. Here the amplitude of oscillation is dependent on the radius from the origin.

**Remark** Phase wrapping is a nice interpretation of shell like features sometimes seen in the outskirts of elliptical galaxies (see <https://apod.nasa.gov/apod/ap140105.html>)

**Remark** Liouville's theorem implies that Hamiltonian phase flows (for 2d phase space) are area preserving.

**Remark** No where did we assume that  $H$  was time independent.






Figure 15: Particles are initially located in the red region but after a period of time are distributed in the green region. Level curves for the dynamical system are circles but the period of oscillation depends on amplitude. The particle distribution is projected downward on the bottom, showing a series of caustics. The process is sometimes described as phase wrapping as the differences in phases of the particles becomes larger with time. The projection is what might be seen in coordinate space. The peaks correspond to extreme positions of radius where the particles spend more time. For example, stars in Keplerian orbit spend more time at apocenter than at pericenter.

## 7 Connections between classical and quantum systems

If the real world is described by a quantum theory, then how is classical physics realized in the limit of  $\hbar \rightarrow 0$ ? Alternatively can classical physics only be realized through *decoherence*? Even though coupled quantum systems would evolve only through unitary evolution, decoherence, which is non-unitary, can be used to describe the behavior of a sub-system. How is classical behavior realized within these pictures? While classical systems are deterministic, measurement in quantum systems involves sampling from a probability distribution. However, the evolution of a density matrix is still deterministic. What types of interesting phenomena arise in these dynamical systems and what are the connections between the different viewpoints?

Given a classical system, what are the different ways to create a related quantum system? How are the properties of the classical system related to the properties of an associated quantum system? What is the connection between integrability of a classical system and integrability of an associated quantum system? For a classical system, ergodicity is the notion that all regions of phase space are frequently visited by a trajectory. What are the properties of the eigenstates of a related quantum system?

Which classical notions of complexity are present in quantum systems and vice versa? What classical notions of complexity are present in measured quantum systems or in quantum systems that are coupled?

We live in a classical world but we can manipulate quantum systems, including quantum computers. Calculations done on quantum computers are likely to live in a hybrid regime where some of the operations are necessarily classical. Does the quantum/classical hybrid regime exhibit different phenomena? How can interesting calculations be carried out in hybrid systems?

### 7.1 Path integrals

In the classical setting, given a Lagrangian  $\mathcal{L}(q, \dot{q}, t)$  and a path  $q(t)$  that goes from  $q_a(t_a)$  to  $q_b(t_b)$ , the action is

$$S[q] \equiv \int_{t_a}^{t_b} \mathcal{L}(q, \dot{q}, t) dt. \quad (46)$$

We are using  $q$  to represent a coordinate and  $[q]$  to represent a particular path. The *principal of least action* for a classical system is that the action  $S[q]$  is minimized over possible paths. In other words, the physically relevant path  $q(t)$  is the one that minimizes the action.

In the quantum setting, instead of minimizing the action, the action is used to construct a transition probability. For a quantum system with Lagrangian  $\mathcal{L}$ , the transition

probability is the modulus of

$$\langle q_a, t_a | q_b, t_b \rangle = \int_{\text{paths}} \mathcal{D}q e^{iS[q]/\hbar} \quad (47)$$

where the integral is not from  $t_a$  to  $t_b$  but over all possible paths between  $q_a$  at time  $t_a$  and  $q_b$  at time  $t_b$ . Here the volume element  $\mathcal{D}q$  represents some way to integrate over possible paths and  $e^{iS[q]/\hbar}$  weights each path with a complex phase. The path integral formulation of quantum field theory (Feynman's path integral formulation) represents the transition amplitude as a weighted sum of all possible histories of the system from the initial to the final state. In the limit  $\hbar \rightarrow 0$ , the weight factor  $e^{iS[q]/\hbar}$  oscillates very rapidly. The main contribution to the path integral comes from paths that make the action stationary and that is consistent with the derivation of the Euler-Lagrange equation from the classical action. Classical physics is realized with the  $\hbar \rightarrow 0$  limit.

## 7.2 Commutation relations

In classical physics, equations of motion can be written in terms of a commutation relation known as a Poisson bracket

$$\{q, p\} = 1. \quad (48)$$

Here  $q, p$  are a coordinate/momentum pair of canonical variables that can be associated with a Hamiltonian. The Poisson bracket of two functions  $f(p, q), g(p, q)$  is more generally

$$\{f, g\} = \frac{\partial f}{\partial q} \frac{\partial g}{\partial p} - \frac{\partial f}{\partial p} \frac{\partial g}{\partial q}. \quad (49)$$

The equations of motion are

$$\dot{q} = \{q, H\} = \frac{\partial H}{\partial p} \quad \dot{p} = \{p, H\} = -\frac{\partial H}{\partial q}. \quad (50)$$

It is tempting to try to *quantize* a system by converting coordinate and momenta into operators  $\hat{x}, \hat{p}$ , and replacing the Poisson bracket with a commutator.

$$[\hat{x}, \hat{p}] = i\hbar \mathbb{I} \quad (51)$$

where  $\mathbb{I}$  is the identity operator. If this worked then equation 50 would be consistent with the Heisenberg version of quantum mechanics where state vectors are fixed but operators evolve, giving for operator  $A(t)$

$$\frac{dA}{dt} = -\frac{i}{\hbar} [A(t), H]. \quad (52)$$

However, it is now known that there is no reasonable quantization map that allows a Poisson bracket to be consistent with a commutator for all functions  $f$  and  $g$ . Work arounds include what is known as *deformation quantization* (which involves a Moyal bracket) and *geometric quantization* which restricts the space of quantizable observables. The Weyl transform (or Weyl quantization) is a way to take a function  $f(p, q)$  and create an operator from it that operates on a Hilbert space. Any observable can be turned into an operator. The inverse transform is known as the Wigner map and generates a function in phase space from an operator.

### 7.3 Quantum chaos

In a classical dynamical system of billiards, a system is defined by a boundary. Inside the boundary (or rink) a particle is free to move on a linear trajectory until it reaches the boundary. At the boundary it bounces, conserving energy. If the rink is 2D and a circle, the dynamics simple to imagine. However, if the rink has bumpers inside or has convex regions, the dynamics can be more complicated and can exhibit chaos in the sense of sensitivity to initial conditions and exponential divergence of nearby orbits. The related quantum system is that of a free wave-like particle within a potential well that becomes infinite at and outside the boundary. The rink can equivalently be described as an acoustic cavity. The presence of classical chaos in the classical billiard system is related to the statistics of the energy spectrum of the equivalent quantum system. This is the setting that is often called *quantum chaos*. Chaos in classical system can also be reflected in how the eigenstates in the quantum system are distributed across the rink.

### 7.4 Decoherence

Beginning with a quantum wavefunction  $|\psi(t)\rangle$  at  $t$ , and a Hamiltonian, its evolution can be specified exactly by Shrodinger's equation;  $i\hbar \frac{d|\psi\rangle}{dt} = H |\psi\rangle$ . This is a dynamical system and it is deterministic. There is no randomness involved. Different experiments give different outcomes only upon measurement (at least within the most common interpretation of quantum mechanics). Decoherence can be described in terms of removing off diagonal elements in a density matrix. However this suggests that the system behaves stochastically, rather than deterministically. How does interaction with another quantum system cause the behavior of a classical system?

### 7.5 Other phenomena

There is a quantum version of the kicked rotor (equivalent to the standard map) that gives rise to Anderson localization.

## 8 Some non-conservative two-dimensional dynamical systems

So far we have only looked at classical Hamiltonian systems (conservative systems). What happens if we add a little bit of damping? The system is not longer area preserving.

### 8.1 Example: Damped pendulum

Consider a damped pendulum model

$$\ddot{\phi} + \alpha \dot{\phi} + \sin \phi = 0 \quad (53)$$

with  $\alpha > 0$ , the damping coefficient. We multiply the equation by  $\dot{\phi}$

$$\begin{aligned} \ddot{\phi} \dot{\phi} + \alpha \dot{\phi}^2 + \dot{\phi} \sin \phi &= 0 \\ \frac{d}{dt} \left[ \frac{\dot{\phi}^2}{2} - \cos \phi \right] &= -\alpha \dot{\phi}^2 \end{aligned}$$

The left term we notice is the energy of the undamped pendulum.

$$\frac{dE}{dt} = -\alpha \dot{\phi}^2$$

This equation holds if we replace  $\cos \phi$  with a general potential  $U(\phi)$ . As long as  $\alpha > 0$  we find that energy is always decreasing. The fixed points of the problem are at same location as for the pendulum ( $\dot{\phi} = 0, \phi = 0, \pi$ ) and their stability is also the same as for the pendulum model. This (and thinking about the energy) makes it straightforward to figure out which direction trajectories go. To help with the phase portrait, we can describe this as a two-dimensional dynamical system with  $\mathbf{x} = (\phi, y)$

$$\begin{aligned} \dot{\phi} &= y \\ \dot{y} &= -\sin \phi - \alpha y \end{aligned} \quad (54)$$

or

$$\begin{pmatrix} \dot{\phi} \\ \dot{y} \end{pmatrix} = \begin{pmatrix} f(y, \phi) \\ g(y, \phi) \end{pmatrix} \quad (55)$$

with  $f(y, \phi) = y$  and  $g(y, \phi) = -\sin \phi - \alpha y$ .

How do we know that this system does not preserve area (is compressible)? Using  $\mathbf{x} = (\phi, y)$  we can compute  $\nabla \cdot \dot{\mathbf{x}}$

$$\begin{aligned} \nabla \cdot \dot{\mathbf{x}} &= \frac{\partial \dot{\phi}}{\partial \phi} + \frac{\partial \dot{y}}{\partial y} \\ &= \frac{\partial f}{\partial \phi} + \frac{\partial g}{\partial y} \\ &= \frac{\partial}{\partial \phi} y + \frac{\partial}{\partial y} (-\sin \phi - \alpha y) \\ &= -\alpha. \end{aligned}$$

This you will recognize is proportional to  $dE/dt$ , the rate that energy is lost. For  $\alpha \neq 0$  the system is not incompressible. Phase space volume is *not* conserved. Our map is *not* an area preserving map.

### 8.1.1 Fixed Points

Consider the Jacobian

$$A = \begin{pmatrix} \frac{\partial f}{\partial \phi} & \frac{\partial f}{\partial y} \\ \frac{\partial g}{\partial \phi} & \frac{\partial g}{\partial y} \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -\cos \phi & -\alpha \end{pmatrix}$$

The Jacobian is useful as near the fixed point we can approximate the dynamics as a linear system with

$$\dot{\mathbf{x}} = A\mathbf{x}$$

where  $\mathbf{x} = (y - y_*, \phi - \phi_*)$  and  $y_*, \phi_*$  are the coordinates of the fixed point. For a linear system a general solution is

$$\mathbf{x} = c_1 e^{\lambda_1 t} \hat{\mathbf{v}}_1 + c_2 e^{\lambda_2 t} \hat{\mathbf{v}}_2$$

where  $\lambda_1, \lambda_2$  are eigenvalues of  $A$ ,  $\mathbf{v}_1, \mathbf{v}_2$  are eigenvectors of  $A$  and  $c_1, c_2$  are constants.

Eigenvalues of the Jacobian matrix at the fixed points can be used to determine the stability of the fixed points. Recall that the trace gives the sum of the eigenvalues and the determinant the product of the two eigenvalues. The characteristic equation can be written

$$\det A - (\text{tr} A)\lambda + \lambda^2 = 0$$

has roots

$$\lambda_{\pm} = \frac{1}{2} \left[ \text{tr} A \pm \sqrt{(\text{tr} A)^2 - 4 \det A} \right]$$

Eigenvalues are both complex (leading to rotation about the fixed point) when

$$(\text{tr} A)^2 - 4 \det A < 0$$

Fixed point is a saddle point if both eigenvalues are real and one is positive and the other negative.

For our damped pendulum  $\text{tr} A = -\alpha$  and  $\det A = \cos \phi$  which is  $\pm 1$  depend on the fixed point.

$$\lambda_{\pm} = \frac{1}{2} \left[ -\alpha \pm \sqrt{\alpha^2 - 4 \cos \phi} \right]$$

At the fixed point with  $\cos \phi = 1$  ( $\phi = 0$ ) we have circulation as long as  $\alpha < 2$ , and in this case both eigenvalues are complex. With  $\alpha > 0$  we expect orbits to spiral in (*stable spiral*). If  $\alpha$  is too large ( $\alpha > 2$ ) then we have two negative eigenvalues, orbits just exponentially drop in without any rotation and the point is just a *stable node*.




Figure 16: Trajectories for a damped pendulum (Equation 53). All trajectories eventually approach a fixed point.

At the other fixed point (with  $\cos \phi = -1$ ,  $\phi = \pi$ ), both eigenvalues are real and one is always positive, the other negative and so it is always a *saddle*.

There are two fixed points, one is always a *saddle* and the other a *sink* (see Figure 16).

### 8.1.2 Appearance of a limit cycle

A slightly more complicated model has the damped pendulum

$$\ddot{\phi} + \alpha \dot{\phi} + \sin \phi = I \quad (56)$$

with constant  $I$ , corresponding to a driven torque. The equations of motion on the plane

$$\begin{aligned} \dot{\phi} &= y \\ \dot{y} &= I - \sin \phi - \alpha y \end{aligned} \quad (57)$$

$$A = \begin{pmatrix} \frac{\partial f}{\partial \phi} & \frac{\partial f}{\partial y} \\ \frac{\partial g}{\partial \phi} & \frac{\partial g}{\partial y} \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -\cos \phi & -\alpha \end{pmatrix}$$

Fixed points are at  $y = 0$ ,  $\sin \phi = I$ , so they only exist when  $|I| < 1$ . We compute  $\text{tr} A = -\alpha$  and  $\det A = \cos \phi$ . At the fixed points  $\det A = \pm \sqrt{1 - I^2}$ . We have a situation similar to above for the damped pendulum where we can have a stable node or a stable spiral for one fixed point where the other is a saddle node.

The fixed point occurs where  $I = \sin \phi$  so if  $I > \pm 1$  then there is no fixed point. However, there could be a stable cycle. For  $I > 1$  we can show that there must be a stable cycle.

Define function  $y_s(\phi)$

$$y_s(\phi) = \frac{1}{\alpha} [I - \sin \phi] \quad (58)$$

where  $\dot{y}$  is zero. Above this line on the  $y, \theta$  cylinder, equation 57 implies that  $\dot{y} < 0$  and below it  $\dot{y} > 0$ . This means that trajectories above and below the  $y_s(\phi)$  curve approach this curve (see Figure 17). All trajectories with  $y$  near  $y_s$  have  $\dot{\phi} > 0$  as  $I > 1$  and  $y_s(\phi) > 0$  (see equations 58 and remembering that  $y = \dot{\phi}$ ) Because  $\dot{\phi} > 0$  all trajectories move to the right on the  $y, \phi$  plane.

Take a point  $y, \phi = 0$  and consider its trajectory when it reaches  $\phi = 2\pi$ . Consider the map  $G(y)$  which takes the initial  $y$  value and returns the  $y$  value at  $\phi = 2\pi$ . A point  $y_a, \phi = 0$  starting above the curve  $y_s$  must have  $G(y_a) < y_a$ . A point  $y_b, \phi = 0$ , starting below the curve  $y_s$  must have  $G(y_b) > y_b$ . At some point the function  $G(y)$  must cross the line  $h(y) = y$  and  $G(y_*) = y_*$  and this orbit corresponds to a periodic orbit (see Figure 18). Since orbits above it move downwards and orbits below it move upwards, it's also a *limit cycle*.

Notes: in regime of high damping  $\alpha > 1$  this system displays hysteresis.  $\langle y \rangle$  is above zero when set by the cycle or equal to zero as set by the fixed point. The dynamical system is used to model a non-linear electronic circuit and Josephson junctions. Limit cycles are examples of phenomena that arise in non-conservative 2 dimensional dynamical systems and not in conservative ones. There are number of ways to prove existence and non-existence of cycles. Also relevant is the Poincaré Bendixon theorem (every limit set of an orbit of a 2d dynamical system is either a fixed point, a periodic orbit or connected set that connects different types of fixed points).




Figure 17: For  $I > 1$  all trajectories approach  $y_s(\phi)$  (equation 58) for the damped torqued pendulum.






Figure 18: The existence of a periodic orbit for the damped torqued pendulum (equation 56).  $G(y_a)$  gives the final  $y$  value of a trajectory that is begun at  $y_a$  and  $\phi = 0$  and integrated until it reaches  $\phi = 2\pi$ . We choose  $y_a$  well below the  $y_s(\phi)$  curve shown in Figure 17 so that  $G(y_a) > y_a$ . We choose  $y_b$  well above the  $y_s$  curve so that  $G(y_b) < y_b$ . Somewhere between  $y_a$  and  $y_b$  there must be a  $y$  value, denoted  $y_*$ , that crosses the  $h(y) = y$  line, giving  $G(y_*) = y_*$ . This is a periodic orbit. Furthermore it is an attracting one, known as a limit cycle.