

## THE VARIABILITY OF SEYFERT 1.8 AND 1.9 GALAXIES AT 1.6 MICRONS

A. C. QUILLEN, SHANNA SHAKED, A. ALONSO-HERRERO, COLLEEN McDONALD, ARIANE LEE, M. J. RIEKE, AND G. H. RIEKE

Steward Observatory, University of Arizona, Tucson, AZ 85721; aquillen@as.arizona.edu

Received 1999 December 9; accepted 2000 February 3; published 2000 March 3

### ABSTRACT

We present a study of Seyfert 1.5–2.0 galaxies observed at two epochs with the *Hubble Space Telescope* (*HST*) at 1.6  $\mu\text{m}$ . We find that unresolved nuclear emission from nine of 14 nuclei varies at the level of 10%–40% on timescales of 0.7–14 months, depending upon the galaxy. A control sample of Seyfert galaxies lacking unresolved sources and galaxies lacking Seyfert nuclei show less than 3% instrumental variation in equivalent aperture measurements. This proves that the unresolved sources are nonstellar and associated with the central parsecs of active galactic nuclei. Unresolved sources in Seyfert 1.8 and 1.9 galaxies are not usually detected in *HST* optical surveys; however, high angular resolution infrared observations will provide a way to measure time delays in these galaxies.

*Subject headings:* galaxies: active — galaxies: nuclei — galaxies: Seyfert — infrared: galaxies

### 1. INTRODUCTION

The discovery that Seyfert 2 galaxies can have reflected or polarized broad line emission has led to an approach coined “unification” toward interpreting the differences between active galactic nuclei (AGNs) in terms of orientation angle (Antonucci 1993). The dusty torus of this unification paradigm absorbs a significant fraction of the optical/UV/X-ray luminosity of an active galaxy and consequently reradiates this energy at infrared wavelengths. As a result of this extinction, it is difficult to observe continuum radiation from Seyfert 2 galaxies at optical and UV wavelengths (e.g., Mulchaey et al. 1994). An additional complication is that in a given aperture it may be difficult to identify the percentage of flux from a nonstellar nuclear source (e.g., Alonso-Herrero, Ward, & Kotilainen 1996). For example, in Seyfert 2 galaxies much of the nuclear emission may originate from nuclear star formation (e.g., Maiolino et al. 1997; Gonzalez-Delgado & Perez 1993).

The high sensitivity and resolution of near-infrared imaging with the *Hubble Space Telescope* (*HST*) using the Near-Infrared Camera and Multiobject Spectrograph (NICMOS) allows us to probe galactic centers at wavelengths that experience reduced extinction compared to the optical and with a beam area about 30 times smaller than is typically achieved with ground-based observations at these wavelengths. This enables us to separate the nuclear emission from that of the surrounding galaxy with unprecedented accuracy. Although a previous survey using WFPC2 at 0.6  $\mu\text{m}$  did not detect unresolved nuclear continuum emission from Seyfert 2 galaxies (Malkan, Gorjian, & Tam 1998), about 60% of the Revised Shapely-Ames and CfA samples (described below) of Seyfert 1.8–2.0 galaxies display prominent unresolved nuclear sources with diffraction rings in NICMOS images at 1.6  $\mu\text{m}$  (McDonald et al. 2000). Although we suspect that these unresolved continuum sources are most likely associated directly with an AGN, they could also be from unresolved star clusters, which are found in a number of normal galaxies (Carollo et al. 1997).

Variability observed in the continuum (e.g., Fitch, Pacholczyk, & Weymann 1967) is an intrinsic property of AGNs which demonstrates that the energy causing the emission must arise from a very small volume. This led early studies to suggest that accretion onto a massive black hole is responsible for the luminosity (Salpeter 1964; Zeldovich & Novikov 1964). Long-term multiyear monitoring programs have found that Sey-

fert 1 galaxies are variable in the near-infrared (Clavel, Wamsteker, & Glass 1989; Lebofsky & Rieke 1980), but these programs have only seen a few Seyfert 2 nuclei vary (e.g., Glass 1997; Lebofsky & Rieke 1980). Evidence for variability in the unresolved sources seen in *HST* observations of Seyfert 2 galaxies would provide evidence that this nuclear emission is nonstellar and so arises from the vicinity of a massive black hole.

### 2. OBSERVATIONS

In this Letter, we present a study of variability in Seyfert galaxies. We have searched the *HST* archive for galaxies (Seyfert and normal) that were imaged *twice* by *HST* at 1.6  $\mu\text{m}$  in the F160W filter with the NICMOS cameras. The Seyfert galaxies with unplanned duplicate observations either satisfy the Revised Shapely-Ames Catalog criteria (described by Maiolino & Rieke 1995) or are part of the CfA redshift survey (Huchra & Burg 1992). The Seyfert observations are discussed in Regan & Mulchaey (1999) and Martini & Pogge (1999), and the observations of the normal or non-Seyfert galaxies are described by Seigar et al. (2000) and Böker et al. (1999). The observations are listed in Table 1 and are grouped by the NICMOS cameras with which they were observed. Images were reduced with the NICRED data reduction software (McLeod 1997) using on orbit darks and flats. Each set of images in the F160W filter was then combined according to the position observed. The pixel sizes for the NICMOS cameras are  $\sim 0''.043$ ,  $0''.076$ , and  $0''.204$  for cameras 1, 2, and 3, respectively.

At the center of these galaxies, we expect contribution from an underlying stellar component in addition to that from an unresolved nonstellar component. To measure the flux from the unresolved component, we must subtract a resolved stellar component. However, this procedure is dependent upon assumptions made about the point-spread function, the form of the stellar surface brightness profile fit to the image, and the region over which we fit this profile. This procedure adds uncertainty in the measurement of the unresolved component. However, aperture photometry has proved quite robust with observations of flux calibration standard stars showing variation less than 1% over the lifetime of NICMOS (M. Rieke 1999, private communication). We therefore opt to use aperture photometry to measure flux variations and then subsequently correct for contamination of the aperture by the background galaxy.

From each pair of images, we measure fluxes in apertures

TABLE 1  
MULTIEPOCH APERTURE PHOTOMETRY OF SEYFERT AND NORMAL GALAXIES AT 1.6  $\mu\text{m}$

GALAXY (1)	LINE CLASSIFICATION (2)	TYPE OF NUCLEUS (3)	FIRST IMAGE		SECOND IMAGE		FLUX		DIFFERENCE (%) (10)
			Proposal ID (4)	Date (5)	Proposal ID (6)	Date (7)	First Image (8)	Second Image (9)	
IC 5063	S2 <sup>a</sup>	D	7330/2	1997 Sep 25	7119/2	1997 Apr 18	2.74	2.73	0.57
NGC 1275	S1.9 <sup>b</sup> /S1.5 <sup>c</sup>	D	7330/2	1998 Mar 16	7457/2	1997 Aug 15	3.74	3.10	19.0
NGC 2460		R	7330/2	1998 Feb 28	7331/2	1997 Sep 11	3.09	3.06	1.06
NGC 2985	T1.9 <sup>c</sup>	R	7330/2	1998 May 18	7331/2	1997 Sep 13	5.44	5.44	-0.04
NGC 3368	L2 <sup>c</sup>	R	7330/2	1998 May 08	7331/2	1998 May 04	9.23	9.33	-1.10
NGC 2903	H <sup>c</sup>	W	7330/2	1998 Apr 22	7331/2	1997 Oct 02	2.96	3.01	-1.66
NGC 6951	S2 <sup>c</sup>	R	7330/2	1998 Mar 30	7331/2	1997 Dec 16	3.88	3.86	0.53
NGC 7177	T2 <sup>c</sup>	R	7330/2	1997 Jun 15	7331/2	1997 Sep 19	2.40	2.46	-2.63
Mrk 266	S2 <sup>b</sup>	R	7867/1	1998 Apr 30	7328/2	1997 Sep 13	1.10	1.07	2.32
Mrk 573	S2 <sup>b</sup>	D	7867/1	1998 Aug 27	7330/2	1997 Jun 26	2.26	2.20	2.69
NGC 3982	S2 <sup>c</sup>	F	7867/1	1998 Sep 10	7330/2	1997 Jun 22	1.36	1.34	1.91
NGC 5033	S1.9 <sup>b</sup> /S1.5 <sup>c</sup>	D	7867/1	1998 Apr 28	7330/2	1997 Aug 19	5.23	6.31	-8.78
NGC 5252	S1.9 <sup>b</sup>	D	7867/1	1998 Mar 09	7330/2	1998 Apr 05	1.81	1.80	0.24
NGC 5273	S1.9 <sup>b</sup> /S1.5 <sup>c</sup>	D	7867/1	1998 May 27	7330/2	1998 Apr 03	2.35	2.57	-8.63
NGC 5347	S2 <sup>d</sup>	D	7867/1	1998 Nov 06	7330/2	1997 Sep 02	2.30	2.18	5.45
NGC 5929	S2 <sup>b</sup>	R	7867/1	1998 Jul 17	7330/2	1998 May 21	1.91	1.90	0.43
Mrk 471	S1.8 <sup>c</sup>	D	7328/1	1998 Apr 16	7867/1	1997 Jul 08	0.88	0.86	2.06
Mrk 533	S2 <sup>b</sup>	D	7328/1	1998 Sep 13	7867/1	1997 Sep 05	4.86	5.34	-9.31
UGC 12138	S1.8 <sup>b</sup>	D	7328/1	1998 Sep 09	7867/1	1997 Jul 28	3.60	4.23	-16.0
UM 146	S1.9 <sup>b</sup>	D	7328/1	1998 Aug 03	7867/1	1997 Sep 13	1.36	1.46	-6.89
NGC 1241	S2 <sup>f</sup>	W	7330/2	1998 Mar 18	7919/3	1998 Jun 19	4.24	4.24	0.01
NGC 214		R	7330/2	1998 May 29	7919/3	1998 Jun 09	2.73	2.67	2.13
NGC 2639	S1.9 <sup>c</sup>	R	7330/2	1998 Feb 23	7919/3	1998 Jun 07	7.25	7.27	-0.29
NGC 2903	H <sup>c</sup>	W	7330/2	1998 Apr 22	7919/3	1998 Jun 09	5.31	5.28	0.45
NGC 3627	T2 <sup>c</sup>	R	7330/2	1998 Apr 22	7919/3	1998 Jun 04	14.99	15.59	-3.89
NGC 404	L2 <sup>c</sup>	D	7330/2	1998 Mar 02	7919/3	1998 Jan 19	11.74	11.64	0.84
NGC 4151	S1.5 <sup>c</sup>	D	7215/2	1998 May 22	7806/3	1997 Oct 14	97.35	108.99	-11.3
NGC 4258	S1.9 <sup>c</sup>	W	7330/2	1997 Nov 21	7919/3	1998 Jun 09	19.20	18.84	1.86
NGC 4395	S1.8 <sup>c</sup>	D	7330/2	1998 May 17	7919/3	1998 Jun 07	1.14	1.02	11.2
NGC 5128	S2 <sup>g</sup>	D	7330/2	1997 Sep 17	7919/3	1998 Jun 17	16.27	16.72	-2.72
NGC 628		W	7330/2	1997 Jun 15	7919/3	1998 Jan 30	1.37	1.40	-2.42
NGC 6744	L <sup>h</sup>	R	7330/2	1997 Sep 09	7919/3	1998 Jun 11	7.03	7.18	-2.08
NGC 6946	H <sup>c</sup>	R	7330/2	1998 May 18	7919/3	1998 Jan 19	13.73	13.99	-1.86

NOTE.—Seyfert and normal galaxies have been grouped by the NICMOS cameras in which they were observed. The first group consists of camera 2/camera 2 pairs, the second camera 1/camera 2 pairs, the third camera 1/camera 1 pairs, and the last group camera 2/camera 3 pairs. Col. (1): Galaxy name. Col. (2): Classification of emission lines in the nucleus. Spectroscopic identifications for the nuclei of NGC 2460 and NGC 214 could not be found. The nucleus of NGC 628 lacks emission lines (Ho et al. 1995). Col. (3): Type of nucleus seen in the F160W images. When the nucleus displayed a clear diffraction ring, it was marked “D,” when the ring was faint it was marked “F” and when the galaxy was resolved, we marked it “R.” When there was an unresolved peak but no sign of a diffraction ring, we marked it “W.” Col. (4): Proposal identification number followed by camera number of the first NICMOS image considered; col. (5): date on which this image was observed. Col. (6): Proposal identification number followed by camera number of the second NICMOS image considered; col. (7): date on which this image was observed. Col. (8): Nuclear flux at 1.6  $\mu\text{m}$  measured in mJy for the image identified by cols. (4) and (5). For the galaxies observed in camera 1 and 2, we used an aperture of 0".602 in diameter. For the galaxies observed solely in camera 1, the aperture was 0".602. For the galaxies observed solely in camera 2, the aperture was 0".760. For the galaxies observed in cameras 2 and 3, the aperture was 1".216. To convert these fluxes into mJy, we used conversion factors  $2.360 \times 10^{-3}$ ,  $2.190 \times 10^{-3}$ ,  $2.776 \times 10^{-3}$  mJy per DN  $\text{s}^{-1}$  for cameras 1, 2, and 3, respectively. This flux calibration is based on measurements of the standard stars P330-E and P172-D during the Servicing Mission Observatory Verification program and subsequent observations (M. Rieke 1999, private communication). Col. (9): Flux in an aperture for the image identified by cols. (6) and (7). Col. (10): Percent difference divided by the mean of the fluxes listed in cols. (8) and (9).

<sup>a</sup> A polarized broad-line component was detected in IC 5063 by Inglis et al. 1993.

<sup>b</sup> From Osterbrock & Martel 1993.

<sup>c</sup> From Ho, Filippenko & Sargent 1995, 1997 (classifications from these works include H = H II nucleus, S = Seyfert nucleus, L = LINER, and T = transition object with numbers corresponding to subtypes).

<sup>d</sup> From Huchra & Burg 1992. No data are available about the line ratios of NGC 5347.

<sup>e</sup> From Dahari, & De Robertis 1988.

<sup>f</sup> From Dahari 1985.

<sup>g</sup> From Tadhunter et al. 1993.

<sup>h</sup> NGC 6744 was classified as a LINER by Vaceli et al. 1997, and no subtype was given.

of the same angular size. No background was subtracted, since the level expected at 1.6  $\mu\text{m}$  is negligible compared to the galaxy surface brightnesses. Apertures are listed in Table 1 and were chosen so that more than 75% of the flux of an unresolved source would be contained in the aperture. We chose apertures based on which two cameras were used to observe the object. We list in Table 1 the difference divided by the mean of the two flux measurements for each pair of images.

To determine whether the nuclear sources are variable, we need to quantify the level of intrinsic scatter in our flux mea-

surements. As a control sample, we use the galaxies not identified as Seyfert galaxies and those containing Seyfert nuclei but lacking an unresolved nuclear component. Comparing camera 2 and camera 3 measurements for this control sample, we find a mean difference of  $\mu = -0.9 \pm 0.7\%$  with a variance of  $\sigma = 2.0\%$  in the measurements. Comparing measurements with two observations in camera 2 for this control sample, we find a mean difference of  $\mu = -0.6 \pm 0.6\%$  with a variance of  $\sigma = 1.4\%$ . Unfortunately, our control sample only contains two galaxies with observations in camera 1 and camera 2 (Mrk 266

TABLE 2  
VARIABLE UNRESOLVED NUCLEAR SOURCES

Seyfert Galaxy (1)	Seyfert Type (2)	$v_{\text{helio}}$ ( $\text{km s}^{-1}$ ) (3)	Size (pc) (4)	Flux (mJy) (5)	Variation (%) (6)	Time (months) (7)	$L_{1.6 \mu\text{m}}$ ( $\text{ergs s}^{-1}$ ) (8)
NGC 1275 <sup>a</sup> .....	S1.9/S1.5	5264	34	3.20	20	7.0	$3.3 \times 10^{42}$
NGC 5033 .....	S1.9/S1.5	875	6	2.41	45	8.3	$7.0 \times 10^{40}$
NGC 5273 .....	S1.9/S1.5	1089	7	1.22	18	1.8	$5.4 \times 10^{40}$
Mrk 533 .....	S2 <sup>b</sup>	8713	56	3.59	13	14.1	$1.0 \times 10^{43}$
NGC 5347 .....	S2	2335	15	1.08	11	12.3	$2.2 \times 10^{41}$
UGC 12138 .....	S1.8	7375	48	2.25	28	13.4	$4.6 \times 10^{42}$
UM 146 .....	S1.9	5208	33	0.60	17	10.7	$6.1 \times 10^{41}$
NGC 4395 <sup>c</sup> .....	S1.8	319	1.3	0.83	15	0.7	$1.2 \times 10^{39}$
NGC 4151 .....	S1.5	995	6.5	103	11	7.3	$3.8 \times 10^{42}$

NOTE.—Seyfert galaxies with unresolved sources that did not vary significantly between observations were IC 5063, Mrk 573, NGC 5252, Mrk 471, and NGC 5128. Col. (3): Heliocentric velocity. Col. (4): Physical size corresponding to  $0''.1$ . These have been estimated using a Hubble constant of  $75 \text{ km s}^{-1} \text{ Mpc}^{-1}$ , except in the case of NGC 4395 for which we adopt a distance of 2.6 Mpc (Rowan-Robinson 1985). Col. (5): The flux of the unresolved component (galaxy subtracted) averaged between the two measurements. We estimate the error to be  $\pm 15\%$  of the flux listed. Col. (6): Percent variation (absolute value of the difference divided by the mean) of the unresolved component. Col. (7): Time between the two different observations. Col. (8): Mean luminosity at  $1.6 \mu\text{m}$  estimated from  $\nu f_{\nu}$ .

<sup>a</sup> Variation in the nuclear flux of NGC 1275 was reported previously by Lebofsky & Rieke 1980.

<sup>b</sup> In Mrk 533, a broad component in Pa $\alpha$  was detected by Ruiz et al. 1994.

<sup>c</sup> NGC 4395 has been labelled “the least luminous Seyfert 1 galaxy” (Filippenko & Sargent 1989; Filippenko, Ho, & Sargent 1993). The flux at  $4400 \text{ \AA}$  has varied by a factor of 3 in just 1 month (Lira et al. 1999), so a variation of 15% in a few weeks at  $1.6 \mu\text{m}$  is not surprising.

and NGC 5929). To supplement this, we also measured stars observed in both camera 1 and 2 in the vicinity of the Galactic center. Differences in fluxes measured in these three image pairs were less than 3%. The statistics of our control sample suggest that the intrinsic scatter of our measurements is smaller than a difference of 3% for all pairs of images. We therefore estimate that flux differences greater than 6% are statistically significant (at  $\geq 2 \sigma$  level) and likely to be caused by variability and not by scatter in the measurements. The galaxies in which we measure differences larger than this level are listed in Table 2.

We did not find that the unresolved nuclear sources in NGC 404 or NGC 2903 were variable. As demonstrated with UV spectra by Maoz et al. (1998), it is possible that the unresolved component in NGC 404 is from a young star cluster. The same is probably true in NGC 2903, which also contains a compact nuclear source and has a nuclear H II region–type spectrum. The scatter in our aperture measurements does not appear to be dependent on the surface brightness profile of the galaxy. No large differences were measured between image pairs for galaxies lacking an unresolved nuclear source.

To estimate the level of variability in the unresolved component, we must measure the contribution within the aperture of this component. For each camera, we measured a point-spread function from stars in the images. We then fit the sum of an exponential bulge profile and the point-spread function to the surface brightness profile. The error in this procedure we estimated from the scatter in the residuals and was about  $\pm 15\%$  of the total unresolved flux measured. We used the flux from the unresolved component and the shape of the point-spread function to estimate the contribution to the flux measured in the apertures listed in Table 1. The differences in the aperture flux measurements are lower limits for differences in the fluxes of the unresolved components (in the limit that the galaxy contributes no flux in these apertures). The mean unresolved fluxes from the two epochs and extent of variability of the unresolved components (the difference divided by the mean) are listed in Table 2.

### 3. DISCUSSION

In nine out of 14 Seyfert 1.5–2.0 galaxies with unresolved components, we find a variation greater than 10% in the flux

of their unresolved continuum nuclear sources in two epochs of observations at  $1.6 \mu\text{m}$ . A control sample of Seyfert galaxies lacking unresolved sources and galaxies lacking Seyfert nuclei show less than 3% instrumental variation in equivalent aperture measurements. This suggests that the variability detected is statistically significant at the level of  $\geq 2 \sigma$ . Since we see variations between 0.7–14 month timescales, the unresolved sources are probably nonstellar and associated with the central parsecs of AGNs. The luminous Seyfert 1 galaxy in our sample, NGC 4151, shows a variation of 10% in its nuclear flux, similar to that seen in the other Seyfert galaxies.

From Table 2 we see that most of the variable sources are Seyfert 1.8 or 1.9 galaxies. NGC 1275, NGC 5033, and NGC 5273 are usually classified as Seyfert 1.9 galaxies, although Ho, Filippenko & Sargent (1995) classify them as S1.5. There are two Seyfert 2 galaxies exhibiting variability: Mrk 533 and NGC 5347. In Mrk 533 a broad component in Pa $\alpha$  was detected by Ruiz, Rieke, & Schmidt (1994), and so this galaxy could be classified as a Seyfert 1.9. Seyfert 1.8 and 1.9 galaxies are more likely to display unresolved nuclear sources than Seyfert 2.0 galaxies (McDonald et al. 2000). In the context of the unification model, reduced extinction toward the continuum-emitting region at  $1.6 \mu\text{m}$  would be expected in Seyfert galaxies that display faint broad-line emission. However, this might also suggest that the sizes of the broad-line region and  $1.6 \mu\text{m}$  continuum emission region are small compared to the material responsible for the bulk of the extinction.

Two major sources for AGN continuum variability are generally considered: (1) instabilities in an accretion disk (e.g., Shakura & Sunyaev 1973) and (2) jet-related processes. The second case could be a possible explanation for variability NGC 1275, since it is bright at radio wavelengths and is significantly polarized at optical wavelengths (as discussed by Angel & Stockman 1980). However, the luminosity of the compact nucleus of this galaxy at 1.3 GHz is about 20 times lower than that we measure at  $1.6 \mu\text{m}$  (using the flux from Taylor & Vermeulen 1996). So the  $1.6 \mu\text{m}$  flux is higher than what would be expected from synchrotron emission and could be from an additional thermal component (e.g., from hot dust). Better measurements showing the shape of the spectral energy distribution spanning the optical and near-infrared region (to see if two

components are present) or a polarization measurement at 1.6  $\mu\text{m}$  would help differentiate between a thermal or non-thermal origin for the near-infrared emission.

For the remainder of the Seyfert galaxies, their low radio power implies that jet-related processes are not responsible for the variability. From observations of the Seyfert 1 galaxy Fairall 9, Clavel et al. (1989) observed large (400 day) time delays between variations seen at 2 and 3  $\mu\text{m}$  and those seen in the UV. Little or no time delay was seen at 1.2  $\mu\text{m}$ . This led them to suggest that the longer wavelength emission was associated with hot dust located outside the broad-line region (e.g., Lebofsky & Rieke 1980; Barvainis 1987; Netzer & Laor 1993) and that the shorter wavelength emission was reprocessed near the UV-emitting region.

For hot dust to cause the 1.6  $\mu\text{m}$  emission, dust grain temperatures resulting from absorption of UV radiation must be quite high, nearly that expected for sublimation ( $T \sim 2000$  K). The grain temperature should reach this level at a radius  $r \sim 0.06 \text{ pc} (L/10^{44} \text{ ergs s}^{-1})^{1/2}$  (following the estimate given in Barvainis 1987). This radius would have a characteristic variability timescale of  $\sim 70$  days or 2 months for a source of  $10^{44} \text{ ergs s}^{-1}$ . We can crudely estimate the bolometric luminosity of our sources from that at 1.6  $\mu\text{m}$  (which are listed in Table 2) by assuming a ratio of  $\sim 10$  between the 1.6  $\mu\text{m}$  and mid-IR luminosity (e.g., Fadda et al. 1998 for the Seyfert 2 galaxies) and a ratio of  $\sim 10$  between the mid-IR and bolometric luminosity (e.g., Spinoglio et al. 1995). The timescales over which we see variations for the brighter sources such as NGC 1275, Mrk 533, and UGC 12138 ( $L \sim 10^{44} \text{ ergs s}^{-1}$ ) are consistent with the 2 month minimum estimated for emission from hot dust. The least luminous of our sources, NGC 4395 ( $L \sim 10^{41} \text{ ergs s}^{-1}$ ), could have a variability timescale of only a few days for hot dust emitting at 1.6  $\mu\text{m}$ , again consistent with the timescale (a few weeks) over which we see a variation.

Emission from hot dust may not necessarily dominate at 1.6  $\mu\text{m}$  since the emitting material would require a temperature near the sublimation point of graphites and silicates (Netzer & Laor 1993). However, transient superheating at larger radii could still cause emission from hot dust at this wavelength. While the timescales over which we see variability are comparable to those expected from hot dust near a sublimation radius, a long-term study comparing flux variations between the near-infrared and X-ray emission would be needed to determine the exact nature of the 1.6  $\mu\text{m}$  emission. This kind of study would also place strong constraints on disk and torus models for the infrared emission (e.g., Efstathiou & Rowan-Robinson 1995; Fadda et al. 1998).

Most of the unresolved nuclear sources studied here exhibit variability. This suggests that most of the many unresolved continuum sources recently discovered in near-infrared surveys (McDonald et al. 2000; Alonso-Herrero et al. 1996)—and not seen in previous optical surveys—are nonstellar and associated with the central parsecs of an AGN. The near-infrared continuum in low-luminosity AGNs can now be studied in a set of objects comprising a larger range of luminosity and orientations. This should provide tests of the unification model for Seyfert 1 and 2 galaxies as well as the nature of accretion in these lower luminosity sources.

We thank the referee, Ski Antonucci, for comments which have improved this Letter. We thank Brad Peterson and Chien Peng for helpful discussions on this work. Support for this work was provided by NASA through grant GO-07869.01-96A from the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Incorporated, under NASA contract NAS5-26555. We also acknowledge support from NASA projects NAG5-3359 and NAG5-3042 and from JPL contract 961633.

#### REFERENCES

- Alonso-Herrero, A., Ward, M. J., & Kotilainen, J. K. 1996, MNRAS, 278, 902  
 Angel, J. R. P., & Stockman, H. S. 1980, ARA&A, 18, 321  
 Antonucci, R. R. J. 1993, ARA&A, 31, 473  
 Barvainis, R. 1987, ApJ, 320, 537  
 Böker, T., et al. 1999, ApJS, 124, 95  
 Carollo, C. M., Stiavelli, M., de Zeeuw, P. T., & Mack, J. 1997, AJ, 114, 2366  
 Clavel, J., Wamsteker, W., & Glass, I. S. 1989, ApJ, 337, 236  
 Dahari, O. 1985, ApJS, 57, 643  
 Dahari, O., & De Robertis, M. M. 1988, ApJS, 67, 249  
 Efstathiou, A., & Rowan-Robinson, M. 1995, MNRAS, 273, 649  
 Fadda, D., Giuricin, G., Granato, G. L., & Vecchies, D. 1998, ApJ, 496, 117  
 Filippenko, A. V., Ho, L. C., & Sargent, W. L. W. 1993, ApJ, 410, L75  
 Filippenko, A. V., & Sargent, W. L. W. 1989, ApJ, 342, L11  
 Fitch, W., Pacholczyk, A. G., & Weymann, R. J. 1967, ApJ, 150, L67  
 Glass, I. S. 1997, MNRAS, 292, L50  
 Gonzalez-Delgado, R. M., & Perez, E. 1993, Ap&SS, 205, 127  
 Ho, L. C., Filippenko, A. V., & Sargent, W. L. W. 1995, ApJS, 98, 477  
 ———. 1997, ApJS, 112, 315  
 Huchra, J., & Burg, R. 1992, ApJ, 393, 90  
 Inglis, M. D., Brindle, C., Hough, J. H., Young, S., Axon, D. J., Bailey, J. A., & Ward, M. J. 1993, MNRAS, 263, 895  
 Lebofsky, M. J., & Rieke, G. H. 1980, Nature, 284, 410  
 Lira, P., Lawrence, A., O'Brien, P., Johnson, R. A., Terlevich, R., & Bannister, N. 1999, MNRAS, 305, 109  
 Maiolino, R., & Rieke, G. H. 1995, ApJ, 454, 95  
 Maiolino, R., Ruiz, M., Rieke, G. H., & Papadopoulos, P. 1997, ApJ, 485, 552  
 Malkan, M. A., Gorjian, V., & Tam, R. 1998, ApJS, 117, 25  
 Maoz, D., Koratkar, A., Shields, J. C., Ho, L. C., Filippenko, A. V., & Sternberg, A. 1998, AJ, 116, 55  
 Martini, P., & Pogge, R. W. 1999, AJ, 118, 2646  
 McDonald, C., Quillen, A. C., Alonso-Herrero, A., Shaked, S., Lee, A., Rieke, M. J., & Rieke, G. H. 2000, ApJ, submitted  
 McLeod, B. 1997, in Proc. 1997 HST Calibration Workshop, ed. S. Casertano, R. Jedrzejewski, T. Keyes, & M. Stevens (Baltimore: STScI), 281  
 Mulchaey, J. S., Koratkar, A., Ward, M. J., Wilson, A. S., Whittle, M., Antonucci, R. R. J., Kinney, A. L., & Hurt, T. 1994, ApJ, 436, 586  
 Netzer, H., & Laor, A. 1993, ApJ, 404, L51  
 Osterbrock, D. E., & Martel, A. 1993, ApJ, 414, 552  
 Regan, M. W., & Mulchaey, J. S. 1999, AJ, 117, 2676  
 Rowan-Robinson, M. 1985, The Cosmological Distance Ladder (New York: Freeman), 171  
 Ruiz, M., Rieke, G. H., & Schmidt, G. D. 1994, ApJ, 423, 608  
 Salpeter, E. E. 1964, ApJ, 140, 796  
 Seigar, M., Stiavelli, M., Carollo, C. M., de Zeeuw, P. T., & DeJongue, H. 2000, in ASP Conf. Ser. 197, Dynamics of Galaxies: From the Early Universe to the Present, ed. F. Combes, G. A. Mamon, & V. Charmandaris (San Francisco: ASP), 269  
 Shakura, N. I., & Sunyaev, R. A. 1973, A&A, 24, 337  
 Spinoglio, L., Malkan, M. A., Rush, B., Carrasco, L., & Recillas-Cruz, E. 1995, ApJ, 453, 616  
 Tadhunter, C. N., Morganti, R., di Serego-Alighieri, S., Fosbury, R. A. E., & Danziger, I. J. 1993, MNRAS, 263, 999  
 Taylor, G. B., & Vermeulen, R. C. 1996, ApJ, 457, L69  
 Vaceli, M. S., Viegas, S. M., Gruenwald, R., & de Souza, R. E. 1997, AJ, 114, 1345  
 Zeldovich, Ya. B., & Novikov, I. D. 1964, Dokl. Akad. Nauk SSSR, 155, 1033